

Polylogarithms at non-positive (i.e. negative) multi-indices

G erard H. E. Duchamp[ ] - Hoang Ngoc Minh[ ] - Ngo Quoc Hoan[ ]

[ ] *Paris XIII University, 93430 Villetaneuse, France, gheduchamp@gmail.com*

[ ] *Lille II University, 59024 Lille, France, Hoang.Ngocminh@lipn.univ-paris13.fr*

[ ] *Paris XIII University, 93430 Villetaneuse, France, quochoan_ngo@yahoo.com.vn*

Abstract

We extend the definition and construct several bases for polylogarithms Li_T , where T is recognizable by a finite state (multiplicity) automaton and of alphabet $X = \{x_0, x_1\}^1$. The kernel of the ‘‘polylogarithmic map’’ Li_\bullet is also characterized and provides a rewriting process which terminates to a normal form. We mostly concentrate on the algebraic aspects of this extension.

- [1] G erard H. E. Duchamp, Hoang Ngoc Minh, Ngo Quoc Hoan, *The algebra of Kleene stars of the plane and polylogarithms*, hal-01267134.

¹The space of rational series considered here is $(\mathbb{C}\langle X \rangle \sqcup \mathbb{C}^{\text{rat}}\langle\langle x_0 \rangle\rangle \sqcup \mathbb{C}^{\text{rat}}\langle\langle x_1 \rangle\rangle, \sqcup, 1_{X^*})$.