

# Polylogarithms at non-positive (i.e. negative) multi-indices

G erard H. E. Duchamp<sup> </sup> - Hoang Ngoc Minh<sup> </sup> - Ngo Quoc Hoan<sup> </sup>

<sup> </sup> *Paris XIII University, 93430 Villetaneuse, France, gheduchamp@gmail.com*

<sup> </sup> *Lille II University, 59024 Lille, France, Hoang.Ngocminh@lipn.univ-paris13.fr*

<sup> </sup> *Paris XIII University, 93430 Villetaneuse, France, quochoan\_ngo@yahoo.com.vn*

---

## Abstract

We extend the definition and construct several bases for polylogarithms  $\text{Li}_T$ , where  $T$  is recognizable by a finite state (multiplicity) automaton and of alphabet  $X = \{x_0, x_1\}^1$ . The kernel of the ‘‘polylogarithmic map’’  $\text{Li}_\bullet$  is also characterized and provides a rewriting process which terminates to a normal form. We mostly concentrate on the algebraic aspects of this extension.

---

- [1] G erard H. E. Duchamp, Hoang Ngoc Minh, Ngo Quoc Hoan, *The algebra of Kleene stars of the plane and polylogarithms*, hal-01267134.

---

<sup>1</sup>The space of rational series considered here is  $(\mathbb{C}\langle X \rangle \sqcup \mathbb{C}^{\text{rat}}\langle\langle x_0 \rangle\rangle \sqcup \mathbb{C}^{\text{rat}}\langle\langle x_1 \rangle\rangle, \sqcup, 1_{X^*})$ .