

# The reflection-absorption model for directed lattice paths

Cyril Banderier<sup>a</sup> Michael Wallner<sup>b</sup>

<sup>a</sup>Laboratoire d'Informatique de Paris Nord, Université Sorbonne Paris Nord, France

<sup>b</sup>Institute of Discrete Mathematics and Geometry, TU Wien, Austria

## Lattice paths

- **Step set:**  $\mathcal{S} = \{s_1, s_2, \dots, s_m\} \subset \mathbb{Z}$
- **$n$ -step lattice path:** sequence of steps  $(v_1, \dots, v_n) \in \mathcal{S}^n$

### Probabilistic weights

- $\Pi = \{p_{s_1}, \dots, p_{s_m}\}$ ,  $p_s \in [0, 1]$  s.t.  $\sum_{s \in \mathcal{S}} p_s = 1$
- **Jump polynomial**  $P(u) = \sum_{s \in \mathcal{S}} p_s u^s$

## The reflection-absorption model

- Lattice:  $\mathbb{Z}_+^2$
- At altitude  $k \neq 0$ 
  - Weighted step set  $\mathcal{S}$
  - $P(u) = \sum_{s \in \mathcal{S}} p_s u^s$
- At altitude  $k = 0$ 
  - Weighted step set  $\mathcal{S}_0$
  - $P_0(u) = \sum_{s \in \mathcal{S}_0} p_{0,s} u^s$




Figure: Allowed steps depend on altitude

**Reflection model:** No loss of mass at 0:  $P_0(1) = 1$

**Absorption model:** Loss of mass at 0:  $P_0(1) < 1$

## Comparing probabilities for different Dyck models

**Step polynomial:**  $P(u) = pu + qu^{-1}$

$P_0(u) =$	bridges, uniform model	absolute value of bridges	excursions, reflection m.	excursions, absorption m.
$pu + qu^{-1}$				
	$\frac{1}{6}$	$\frac{1}{1+p_0/p+q_0/q}$	$\frac{p}{1+p}$	$\frac{p}{p+p_0}$
	$\frac{1}{6}$	$\frac{p_0/p+q_0/q}{1+p_0/p+q_0/q}$	$\frac{1}{1+p}$	$\frac{p_0}{p+p_0}$
	$\frac{1}{6}$	0	0	0
	$\frac{1}{6}$	0	0	0
	$\frac{1}{6}$	0	0	0
	$\frac{1}{6}$	0	0	0

## Relevant constants

Let  $\tau$  be the *structural constant* given by  $P'(\tau) = 0$ ,  $\tau > 0$ , and let  $\rho = 1/P(\tau)$  be the *structural radius*.

Let  $u_1(z)$  be the unique solution of the *kernel equation*

$$1 - zP(u) = 0,$$

with  $\lim_{z \rightarrow 0} u_1(z) = 0$ . Then, the equation  $1 - zP_0(u_1(z)) = 0$  has at most one solution in  $z \in (0, \rho]$ , which we denote by  $\rho_1$ .

Additionally, we define the constants

$$\alpha = (P_0(u_1(z)))'|_{z=\rho_1}, \quad \alpha_2 = (P_0(u_1(z)))''|_{z=\rho_1},$$

$$\gamma = \frac{1}{\alpha\rho_1^2 + 1}, \quad \kappa = \rho\sqrt{2\frac{P(\tau)}{P''(\tau)}P_0'(\tau)}, \quad \lambda = \frac{P_0(\tau)}{P(\tau)}.$$

## Asymptotic number of excursions

Let  $e_n$  be number of excursions of length  $n$ . Then, the generating function of excursions is of the kind

$$E(z) := \sum_{n \geq 0} e_n z^n = \frac{1}{1 - zP_0(u_1(z))}.$$

The asymptotic number of excursions is given by

$$e_n \sim \begin{cases} \gamma\rho_1^{-n} (1 + \mathcal{O}(\frac{1}{n})), & \text{supercritical case: } \lambda > 1, \\ \frac{1}{\kappa\sqrt{\pi n^{1/2}}} (1 + \mathcal{O}(\frac{1}{n})), & \text{critical case: } \lambda = 1, \\ \frac{\kappa}{2\sqrt{\pi(1-\rho P_0(\tau))^2 n^{3/2}}} (1 + \mathcal{O}(\frac{1}{n})), & \text{subcritical case: } \lambda < 1. \end{cases}$$

## Returns to zero

### Definition

- **Arch:** excursion of size  $> 0$  whose only contacts with the  $x$ -axis are at its end points.
- **Return to zero:** vertex of a path of altitude 0 with positive abscissa.




Figure: An excursion with 3 returns to zero

### Corresponding generating function

$$E(z, u) := \sum_{n, k \geq 0} e_{n,k} z^n u^k = \frac{1}{1 - uzP_0(u_1(z))}$$

### Excursion of length $n$ having $k$ returns to zero

$$\mathbb{P}(X_n = k) := \mathbb{P}(\text{size} = n, \# \text{returns to zero} = k) = \frac{e_{n,k}}{e_n}$$


## Limit laws for returns to zero of excursions

1. In the supercritical case, i.e.  $\lambda > 1$ ,

$$\frac{X_n - \mu n}{\sigma\sqrt{n}}, \quad \text{with}$$


$$\mu = \gamma, \quad \sigma = \gamma^3(\alpha_2\rho_1^3 - 2) + \gamma^2(\rho_1 + 2) - \gamma,$$

converges in law to a **Gaussian variable**  $N(0, 1)$ .




2. In the critical case, i.e.  $\lambda = 1$ , the normalized random variable  $\frac{\kappa}{\sqrt{2n}}X_n$ , converges in law to a **Rayleigh distribution** (density:  $xe^{-x^2/2}$ ):

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\frac{\kappa}{\sqrt{2n}}X_n \leq x\right) = 1 - e^{-x^2/2}.$$



3. In the subcritical case, i.e.  $\lambda < 1$ , the limit distribution of  $X_n - 1$  is the **negative binomial distribution**  $\text{NegBin}(2, 1 - \lambda)$ :

$$\mathbb{P}(X_n - 1 = k) \sim (k + 1)\lambda^k(1 - \lambda)^2.$$



## Conclusions

- Similar results hold for the asymptotics of bridges and meanders,
- Limit laws for other parameters like final altitude of meanders, or returns to zero of bridges exist,
- Extensions to more general lattice path models are possible.

## References

- [1] Banderier, C.; Flajolet, P.: *Basic analytic combinatorics of directed lattice paths*. Theoretical Computer Science, 281, p. 37-80, 2002.
- [2] Banderier, C.; Wallner, M.: *Some reflections on directed lattice paths*. Proceedings of the 25th Int. Conf. on Prob., Comb. and Asymptotic Methods for the Analysis of Algorithms (AofA'14), p. 25-36, 2014.
- [3] Flajolet, P.; Sedgewick, R.: *Analytic Combinatorics*. Cambridge University Press, 2009.