

# A Three Parameter Hopf Deformation of the Algebra of Feynman-like Diagrams

G H E Duchamp<sup>a</sup>, K A Penson<sup>c</sup>, A I Solomon<sup>c,d</sup>, P Blasiak<sup>b</sup>  
and A Horzela<sup>b</sup>

<sup>a</sup> Institut Galilée, LIPN, CNRS UMR 7030  
99 Av. J.-B. Clement, F-93430 Villetaneuse, France

<sup>b</sup> H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences  
ul. Eliasza-Radzikowskiego 152, PL 31342 Kraków, Poland

<sup>c</sup> Laboratoire de Physique Théorique de la Matière Condensée  
Université Pierre et Marie Curie, CNRS UMR 7600  
Tour 24 - 2ième ét., 4 pl. Jussieu, F 75252 Paris Cedex 05, France

<sup>e</sup> The Open University, Physics and Astronomy Department  
Milton Keynes MK7 6AA, United Kingdom

E-mail:

ghed@lipn-univ.paris13.fr, a.i.solomon@open.ac.uk  
pawel.blasiak@ifj.edu.pl, andrzej.horzela@ifj.edu.pl,  
penson@lptl.jussieu.fr,

**Abstract.** We construct in this work a three parameter deformation of the Hopf algebra (that we call **LDIAG**) appearing in an expansion, in terms of Feynman-like Diagrams, of the product formula in simplified version of Quantum Field Theory. This new algebra is a true Hopf deformation which specializes to **LDIAG** for some parameter values and to the algebra of Matrix Quasi-Symmetric Functions (**MQSym**) for other values, relating **LDIAG** to other Hopf algebras of contemporary Physics. Moreover there is an onto linear mapping preserving products from our algebra to the algebra of Euler-Zagier sums.

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>How and why these Feynman-like Diagrams arise</b>	<b>3</b>
<b>3</b>	<b>Non-commutative lifting (classical case)</b>	<b>6</b>
3.1	Free monoids . . . . .	6
3.2	Labelling the nodes . . . . .	7
3.3	Coding <b>ldiag</b> with “lists of monomials” . . . . .	8

<i>CONTENTS</i>	2
<b>4 The Hopf algebra LDIAG (non-deformed case)</b>	<b>8</b>
4.1 The monoid $(\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(X))^*$ and the submonoid of codes of diagrams . . .	9
4.2 The Hopf algebras $\mathcal{C}_{ldiag}$ and <b>LDIAG</b> . . . . .	10
4.3 Subalgebras of <b>LDIAG</b> . . . . .	13
4.3.1 Graphic primitive elements . . . . .	13
4.3.2 Level subalgebras . . . . .	14
4.3.3 <b>BELL</b> and <b>LBELL</b> . . . . .	14
<b>5 The algebra LDIAG(<math>q_c, q_s, q_t</math>) (deformed case)</b>	<b>15</b>
5.1 Counting crossings ( $q_c$ ) and superpositions ( $q_s$ ) . . . . .	15
5.2 Modified laws . . . . .	16
<b>6 Coproducts</b>	<b>19</b>
<b>7 LDIAG(<math>q_c, q_s, q_t</math>) and its specializations</b>	<b>21</b>
<b>8 Conclusion</b>	<b>21</b>

## 1. Introduction

Let us describe roughly the route from the product formula by Bender et al. [3] and their related Feynman-like diagrams to the discovery of Hopf algebra structures [14] on the diagrams themselves compatible with their evaluations.

First, C. M. Bender, D. C. Brody, and B. K. Meister [3] introduced a special field theory which proved to be particularly rich in combinatorial links and by-products.

Second, the Feynman-like diagrams produced by this theory label monomials and we had the surprise to see that they naturally combine in a way compatible with the monomial multiplication and co-addition. This is the Hopf algebra **DIAG**.

Third, the natural noncommutative pull-back of this algebra, **LDIAG**, has a basis (the labelled diagrams) which is in one-to-one correspondence with that of the Matrix Quasi-Symmetric Fonctions (the packed matrices of **MQSym**), but their algebra and co-algebra structures are completely different. In particular, in this basis, the multiplication of **MQSym** implies a sort of shifted shuffle with overlappings reminiscent to Hoffmann's shuffle used in the theory of polyzeta functions [11]. The superpositions and overlappings involved there are not present in the (non-deformed) **LDIAG** and, moreover, the coproduct of **LDIAG** is co-commutative and the one of **MQSym** is not.

The aim of this paper is to introduce some “parametric algebra” between the two Hopf algebras **LDIAG** and **MQSym**. The striking result is that, introducing parameters to count, in the most natural way, the crossings and overlappings of the shifted shuffle one can witness that the resulting law is associative (graded with unit). We also show how to interpolate with a coproduct which makes, at each stage, our algebra a Hopf algebra. The result is thus a three-parameters Hopf algebra deformation which specialises to **LDIAG** at  $(0, 0, 0)$  and to **MQSym** at  $(1, 1, 1)$ . Moreover it appears that, for some set of parameters, the multiplication rule of **LDIAG** covers the one of Euler-Zagier sums.

ACKNOWLEDGEMENTS : The first material of this paper was set during the staying of one of the authors at “the first” (Exeter). Be its owner, Catherine Borgen, be thanked for her creative hospitality.

The structure of the paper is the following ...

## 2. How and why these Feynman-like Diagrams arise

The beginning of the story was explained with full details in [33, 34, 35, 26, 5, 6], and the Hopf algebra structure is made precise in [14, 37]. Here, we will make the beginning shorter but focus on the last part, where the algebraic structure constructed on the diagrams themselves arise.

The very starting point is the formula (product formula) of Bender and al. [3], which can be considered as an expression of the Hadamard product for exponential generating series. That is, with

$$F(z) = \sum_{n \geq 0} a_n \frac{z^n}{n!}, \quad G(z) = \sum_{n \geq 0} b_n \frac{z^n}{n!}, \quad \mathcal{H}(F, G) := \sum_{n \geq 0} a_n b_n \frac{z^n}{n!} \quad (1)$$

one can check that

$$\mathcal{H}(F, G) = F \left( z \frac{d}{dx} \right) G(x) \Big|_{x=0}. \quad (2)$$

This product being bilinear, in the case when  $F(0)$  and  $G(0)$  are not zero, one can renormalize the functions so that  $F(0) = G(0) = 1$  and one is interested to obtain compact and generic formulas. If we write the functions as

$$F(z) = \exp \left( \sum_{n=1}^{\infty} L_n \frac{z^n}{n!} \right), \quad G(z) = \exp \left( \sum_{n=1}^{\infty} V_n \frac{z^n}{n!} \right). \quad (3)$$

that is as free exponentials, one gets, through the Bell polynomials in the sets of variables  $\mathbb{L}, \mathbb{V}$  (see [14] for details)

$$\mathcal{H}(F, G) = \sum_{n \geq 0} \frac{z^n}{n!} \sum_{P_1, P_2 \in UP_n} \mathbb{L}^{Type(P_1)} \mathbb{V}^{Type(P_2)} \quad (4)$$

where  $UP_n$  is the set of unordered partitions of  $[1 \cdots n]$ . An unordered partition  $P$  of a set  $X$ , is a subset of  $P \subset \mathfrak{P}(X) - \{\emptyset\}^\dagger$  (that is an unordered collection of blocks, i. e. non-empty subsets of  $X$ ) such that

- the union  $\bigcup_{Y \in P} Y = X$  ( $P$  is a covering)
- $P$  consists of disjoint subsets, i. e.  
 $Y_1, Y_2 \in P$  and  $Y_1 \cap Y_2 \neq \emptyset \implies Y_1 = Y_2$ .

The type of  $P \in UP_n$  (denoted above  $Type(P)$ ) is the multiindex  $(\alpha_i)_{i \in \mathbb{N}^+}$  such that  $\alpha_k$  is the number of  $k$ -blocks, that is the number of members of  $P$  with cardinality  $k$ .

Here is the point where the formula entangles and the diagrams of the theory arise. The fundamental remarks are :


- the monomial  $\mathbb{L}^{Type(P_1)} \mathbb{V}^{Type(P_2)}$  needs much less information than that which is contained in the individual partitions  $P_1, P_2$  (for example, one can relabel the elements without changing the monomial),
- two partitions have an incidence matrix *from which it is still possible to recover the types of the partitions*

Now, the construction goes as follows.

† The set of subsets of  $X$  is denoted  $\mathfrak{P}(X)$  (this denotation [9] owes to the former german school).

- (i) Take two unordered partitions of  $[1 \cdots n]$ , say  $P_1, P_2$
- (ii) Build their incidence matrix  $(\text{card}(Y \cap Z))_{(Y,Z) \in P_1 \times P_2}$
- (iii) Build the diagram representing the multiplicities of the incidence matrix : for each block of  $P_1$  draw a black spot (resp. for each block of  $P_2$  draw a white spot)
- (iv) Draw lines between the black spot  $Y \in P_1$  and the white spot  $Z \in P_2$  as much as  $\text{card}(Y \cap Z)$
- (v) Remove the information of the blocks  $Y, Z, \dots$ .

So doing, one obtains a bipartite graph with  $p$  ( $= \text{card}(P_1)$ ) black spots,  $q$  ( $= \text{card}(P_2)$ ) white spots, no isolated vertex and integer multiplicities. Their set will be denoted **diag**.



**Fig 1.** — Diagram from  $P_1, P_2$  (set partitions of  $[1 \cdots 11]$ ).

$P_1 = \{\{2, 3, 5\}, \{1, 4, 6, 7, 8\}, \{9, 10, 11\}\}$  and  $P_2 = \{\{1\}, \{2, 3, 4\}, \{5, 6, 7, 8, 9\}, \{10, 11\}\}$  (respectively black spots for  $P_1$  and white spots for  $P_2$ ).

The incidence matrix corresponding to the diagram (as drawn) or these partitions is  $\begin{pmatrix} 0 & 2 & 1 & 0 \\ 1 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \end{pmatrix}$ . But, due to the fact that the defining partitions are unordered, one can permute the spots (black and white, between themselves) and, so, the lines and columns of this matrix can be permuted. the diagram could be represented by the matrix  $\begin{pmatrix} 0 & 0 & 1 & 2 \\ 0 & 2 & 1 & 0 \\ 1 & 0 & 3 & 1 \end{pmatrix}$  as well.

The product formula now reads

$$\mathcal{H}(F, G) = \sum_{n \geq 0} \frac{z^n}{n!} \sum_{\substack{d \in \text{diag} \\ |d|=n}} \text{mult}(d) \mathbb{L}^{\alpha(d)} \mathbb{V}^{\beta(d)} \tag{5}$$

where  $\alpha(d)$  (resp.  $\beta(d)$ ) is the “white spots type” (resp. the “black spots type”) i.e. the multiindex  $(\alpha_i)_{i \in \mathbb{N}^+}$  (resp.  $(\beta_i)_{i \in \mathbb{N}^+}$ ) such that  $\alpha_i$  (resp.  $\beta_i$ ) is the number of white spots (resp. black spots) of degree  $i$  ( $i$  lines connected to the spot) and  $\text{mult}(d)$  is the number of pairs of unordered partitions of  $[1 \cdots |d|]$  (here  $|d| = |\alpha(d)| = |\beta(d)|$  is the number of lines of  $d$ ) with associated diagram  $d$ .

Now the natural question arises :

Q1) “Is there a (graphically) natural multiplicative structure on **diag** such that the arrow

$$d \mapsto \mathbb{L}^{\alpha(d)} \mathbb{V}^{\beta(d)} \tag{6}$$

be a morphism ?”

The answer is “yes”. The desired product just consists in concatenating the diagrams (the result, i. e. the diagram obtained in placing  $d_2$  at the right of  $d_1$ , will be denoted  $[d_1|d_2]_D$ ). One must check that this product is compatible with the permutation (of white and black spots between themselves) equivalence, which is rather straightforward (see [14]). We have

**Proposition 2.1** *Let  $\mathbf{diag}$  be the set of diagrams (including the void one).*

*i) The law  $(d_1, d_2) \mapsto [d_1|d_2]_D$  endows  $\mathbf{diag}$  with the structure of a commutative monoid with the void diagram as neutral (this diagram will, therefore, be denoted  $1_{\mathbf{diag}}$ ).*

*ii) The arrow  $d \mapsto \mathbb{L}^{\alpha(d)}\mathbb{V}^{\beta(d)}$  is a morphism of monoids, the codomain of this arrow being the monoid of (commutative) monomials in the alphabet  $\mathbb{L} \cup \mathbb{V}$  i.e.*

$$\mathfrak{M}\mathfrak{O}\mathfrak{N}(\mathbb{L} \cup \mathbb{V}) = \{\mathbb{L}^{\alpha}\mathbb{V}^{\beta}\}_{\alpha, \beta \in (\mathbb{N}^+)^{(\mathbb{N})}} = \bigcup_{n, m \geq 1} \{L_1^{\alpha_1} L_2^{\alpha_2} \cdots L_n^{\alpha_n} V_1^{\beta_1} V_2^{\beta_2} \cdots V_m^{\beta_m}\}_{\alpha_i, \beta_j \in \mathbb{N}}.$$

*iii) The monoid  $(\mathbf{diag}, [-|-]_D, 1_{\mathbf{diag}})$  is a free commutative monoid. Its letters are the connected (non void) diagrams.*

**Remark 2.2** *The reader which is not familiar with the algebraic structure of  $\mathfrak{M}\mathfrak{O}\mathfrak{N}(\mathbb{X})$  can find rigorous definitions in paragraph (3.1) where this structure is needed to prepare the proofs relating to deformations.*

### 3. Non-commutative lifting (classical case)

The “classical” construction of the Hopf algebra **LDIAG** was announced in [14]. We give below the proofs using a coding through “lists of monomials” which prepares the deformed (quantum) case. The entries of a list can be considered as “coordinate functions” for the diagrams (see introduction of section (4)).

#### 3.1. Free monoids

We here recall the construction of the free and free commutative monoids generated by a given set of variables (i.e. an alphabet).

Let  $\mathbb{X}$ , be a set. We denote by  $\mathbb{X}^*$  the set of lists of elements of  $\mathbb{X}$ , including the void one. In many works, and in the sequel, the list  $[x_1, x_2, \cdots, x_n]$  will be denoted as a word  $x_1x_2 \cdots x_n$  so that the concatenation of two lists  $[x_1, x_2, \cdots, x_n]$ ,  $[y_1, y_2, \cdots, y_m]$  is just the word  $x_1x_2 \cdots x_ny_1y_2 \cdots y_m$ . For this (associative) law, the void list  $[\ ]$  is neutral and will therefore be denoted  $1_{\mathbb{X}^*}$

Similarly, we denote  $\mathbb{N}^{(\mathbb{X})}$  [7] the set of multisubsets of  $\mathbb{X}$  (i.e. the set of - multiplicity - mappings with finite support  $\mathbb{X} \mapsto \mathbb{N}$ ). Every element  $\alpha$  of  $\mathbb{N}^{(\mathbb{X})}$  can be written multiplicatively, following the classical multiindex notation

$$\mathbb{X}^{\alpha} = \prod_{x \in \mathbb{X}} x^{\alpha(x)} \tag{7}$$


and the set  $\mathfrak{M}\mathfrak{D}\mathfrak{N}(X) = \{\mathbb{X}^\alpha\}_{\alpha \in \mathbb{N}(X)}$  is exactly the set of (commutative) monomials with variables in  $\mathbb{X}$ . It is a monoid, indeed a (multiplicative) copy of  $\mathbb{N}(X)$  as  $\mathbb{X}^\alpha \mathbb{X}^\beta = \mathbb{X}^{\alpha+\beta}$ . The subset of its non-unit elements is a semigroup which will be denoted by  $\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(X)$  ( $= \mathfrak{M}\mathfrak{D}\mathfrak{N}(X) - \{\mathbb{X}^0\}$ ).

### 3.2. Labelling the nodes

There are (at least) two good reasons to look for non-commutative structures which may serve as a noncommutative pullback for **diag**.

- Rows and Columns of matrices are usually (linearly) ordered and we have seen that a diagram is not represented by a matrix but by a class of matrices
- The “expressive power” of **diag** and its algebra is not sufficient to connect it to other (non-commutative or non-cocommutative) algebras relevant in contemporary physics

The solution (of the non-deformed problem [14]) is simple and consists in labelling the nodes from left to right and from “1” to the desired amount as follows.



**Fig 2.** — *Labelled diagram of format  $3 \times 4$  corresponding to the one of Fig 1.*

The set of these graphs (i.e. bipartite graphs on some product  $[1..p] \times [1..q]$  with no isolated vertex) will be denoted **ldiag**. The composition law is, as previously, the concatenation in the obvious sense. Explicitly, if  $d_i$ ,  $i = 1, 2$  are two diagrams of dimensions  $[1..p_i] \times [1..q_i]$ , one relabels the black (resp. white) spots of  $d_2$  from  $p_1 + 1$  to  $p_1 + p_2$  (resp. from  $q_1 + 1$  to  $q_1 + q_2$ ) the result will be noted  $[d_1|d_2]_L$ . One has

**Proposition 3.1** *Let **ldiag** be the set of labelled diagrams (including the void one).*

*i) The law  $(d_1, d_2) \mapsto [d_1|d_2]_L$  endows **ldiag** with a structure of noncommutative monoid with the void diagram ( $p = q = 0$ ) as neutral (which will, therefore, be denoted  $1_{\mathbf{ldiag}}$ ).*

*ii) The arrow from **ldiag** to **diag**, which consists in “forgetting the labels of the vertices” is a morphism of monoids.*

*iii) The monoid  $(\mathbf{ldiag}, [-|-]_L, 1_{\mathbf{ldiag}})$  is a free (noncommutative) monoid. Its letters are the irreducible diagrams (denoted, from now on  $\mathit{irr}(\mathbf{ldiag})$ ).*


**Remark 3.2** *i) In a general monoid  $(M, \star, 1_M)$ , the irreducible elements are the elements  $x \neq 1_M$  such that  $x = y \star z \implies 1_M \in \{y, z\}$ .*

*ii) It can happen that an irreducible of **ldiag** has an image in **diag** which splits as shows the simple example of the cross defined by the incidence matrix  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .*

### 3.3. Coding **ldiag** with “lists of monomials”

One can code every labelled diagram by a “list of (commutative) monomials” in the following way.

- Let  $\mathbb{X} = \{x_i\}_{i \geq 1}$  an infinite set of indeterminates and  $d \in \mathbf{ldiag}_{p \times q}$  a diagram ( $\mathbf{ldiag}_{p \times q}$  is the set of diagrams with  $p$  black spots and  $q$  white spots).
- Associate to  $d$  the multiplicity function  $[1..p] \times [1..q] \rightarrow \mathbb{N}$  such that  $d(i, j)$  is the number of lines from the black spot  $i$  to the white spot  $j$ .
- The code associated to  $d$  is  $\varphi_{lm}(d) = [m_1, m_2, \dots, m_p]$  such that  $m_i = \prod_{j=1}^q x_j^{m(i,j)}$



**Fig 3.** — Coding the diagram of fig 2 by a word of monomials. The code here is  $[x_2^2 x_3, x_1 x_2 x_3^3, x_3 x_4^2]$

As a data structure, the lists of monomials are elements of  $(\mathfrak{MON}^+(X))^*$ , the free monoid whose letters are  $\mathfrak{MON}^+(X) = \mathfrak{MON}(X) - \{\mathbb{X}^0\}$ , the semigroup of non-unit monomials over  $\mathbb{X}$ .

It is not difficult to see that, through this coding, the concatenation reflects according to the following formula

$$\varphi_{lm}([d_1 | d_2]_L) = \varphi_{lm}(d_1) * T_{\max(\text{IndAlph}(\varphi_{lm}(l_1)))}(\varphi_{lm}(d_2)) \quad (8)$$

where  $T_p$  is the translation operator which changes the variables according to  $T_p(x_i) = x_{i+p}$  (which corresponds to the relabelling of the white spots) and  $p_1$  is the number of black spots of  $d_1$ .

For example, one has

$$T_2([x_2^2 x_3, x_1 x_2 x_3^3, x_3 x_4^2]) = [x_4^2 x_5, x_3 x_4 x_5^3, x_5 x_6^2] ; T_6([x_1, x_2^2]) = [x_7, x_8^2] \quad (9)$$

## 4. The Hopf algebra LDIAG (non-deformed case)

In [14], we constructed a structure of Hopf algebra structure on the space of diagrams **LDIAG**. The aim of this section is to give complete proofs and details for this



construction through the use of the special space of coordinates constructed above (the complete vector of coordinates of a diagram being its code).

4.1. The monoid  $(\mathfrak{MON}^+(X))^*$  and the submonoid of codes of diagrams

Formula (8) can be written with the lists

$$l_1 \bar{*} l_2 = l_1 * T_{\max(\text{IndAlph}(l_1))}(l_2) \quad (10)$$

defines a structure of monoid on  $(\mathfrak{MON}^+(X))^*$  (the set of lists of non-unit monomials) with the empty list as neutral (i.e.  $[\ ]$  which will, therefore, be denoted by “ $1_{(\mathfrak{MON}^+(X))^*}$ ” or simply “1” when the context is clear).

We will return to this construction (called shifting [19]) later.

The alphabet of a list is the set of variables occurring in the list. Formally

$$\text{Alph}([m_1, m_2, \dots, m_n]) = \bigcup_{1 \leq i \leq n} \text{Alph}(m_i) \quad (11)$$

where, classically, for a monomial  $m = \mathbb{X}^\alpha$ ,  $\text{Alph}(m) = \{x_i\}_{\alpha(i) \neq 0}$ .

Now, we can define the “compacting operator” on  $k\langle \mathfrak{MON}^+(X) \rangle$  by its action on the lists. This operator actually removes the holes in the alphabet of a list by pushing to the left the indices which are at the right of a hole. For example (we denote  $cpt$  the operator)

$$cpt([x_2^2 x_{10}, x_3 x_4 x_8^3, x_3 x_4^2]) = [x_1^2 x_5, x_2 x_3 x_4^3, x_2 x_3^2] \quad (12)$$

the alphabet of the list in the LHS is  $\text{Alph}(l) = \text{Alph}([x_2^2 x_{10}, x_3 x_4 x_8^3, x_3 x_4^2]) = \{x_2, x_3, x_4, x_8, x_{10}\}$ , its indices are  $\text{IndAlph}(l) = \{2, 3, 4, 8, 10\}$  and the reindexing function is the unique strictly increasing mapping from  $\{2, 3, 4, 8, 10\}$  to  $[[5]]$ . The compacting operator is, here, just the substitution

$$x_1 \leftarrow x_2; x_2 \leftarrow x_3; x_3 \leftarrow x_4; x_4 \leftarrow x_8; x_5 \leftarrow x_{10}$$

The formal definitions are the following

- $\text{IndAlph}(l) = \{i \mid x_i \in \text{Alph}(l)\}$
- $l$  being given, let  $\phi_l$  be the unique increasing mapping from  $\text{IndAlph}(l)$  to  $[[\text{card}(\text{IndAlph}(l))]]$  (in fact,  $\text{card}(\text{IndAlph}(l)) = \text{card}(\text{Alph}(l))$ )
- let  $s_l$  be the substitution  $x_i \leftarrow x_{\phi_l(i)}$  in the monomials.
- Then, if  $l = [m_1, m_2, \dots, m_n]$ ,  $cpt(l) = [s_l(m_1), s_l(m_2), \dots, s_l(m_n)]$ .

**Définition 4.1** *The compacting operator  $cpt : k\langle \mathfrak{MON}^+(X) \rangle \mapsto k\langle \mathfrak{MON}^+(X) \rangle$  is the extension by linearity of the mapping  $cpt$  defined above.*

It can be checked easily that, for  $l \in (\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(X))^*$ , the following are equivalent

- (i)  $cpt(l) = l$
- (ii)  $IndAlph(l) = [[card(IndAlph(l))]]$
- (iii) there is no hole in  $Alph(l)$  that is there exists no  $i \geq 1$  s.t.  $x_i \notin Alph(l)$  and  $x_{i+1} \in Alph(l)$
- (iv)  $l$  is the code of some (then unique) diagram  $d$ .

It follows from the preceding properties that  $cpt$  is a projector with range the subspace  $\mathcal{C}_{ldiag}$  of  $k\langle\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X})\rangle$  generated by the codes of the diagrams. Formula (8) proves that  $\mathcal{C}_{ldiag}$  is closed under the shifted concatenation defined by (10). More precisely

**Proposition 4.2** *The algebra  $\mathcal{C}_{ldiag}$  is a free algebra on the set of the codes of irreducible diagrams.*

These codes are also the non-empty lists  $l$  which are compact (i.e.  $cpt(l) = l$ ) and cannot be factorized into a product of two non-empty lists i.e.  $l = l_1 * l_2$ ;  $l_i \neq [ ]$  (one can check easily that, if  $l_1 * l_2$  is compact, so are  $l_1$  and  $l_2$ ).

#### 4.2. The Hopf algebras $\mathcal{C}_{ldiag}$ and **LDIAG**

The algebra **LDIAG** is endowed with a structure of a bi-algebra by the comultiplication

$$\Delta_{BS}(d) = \sum_{I+J=[1..p]} d[I] \otimes d[J] \quad (13)$$

where  $p$  is the number of black spots and  $d[I]$  is the “restriction” of  $d$  to the black spots selected by  $I \subset [1..p]$ .

On the other hand, we have the standard Hopf algebra structure on the Free algebra, expressed in terms of concatenation and subwords [24, 31]. Let  $\mathbb{A}$  be an alphabet (a set of letters) and  $w \in \mathbb{A}^*$  a word, if we write  $w$  a sequence of letters  $w = a_1 a_2 \cdots a_n$ ;  $a_i \in \mathbb{A}$ , the length  $|w|$  of  $w$  is  $n$  and if  $I = \{i_1, i_2, \cdots, i_k\} \subset [1..n]$ , the subword  $w[I]$  is  $a_{i_1} a_{i_2} \cdots a_{i_k}$  (this notation is slightly different from that of [31] where it is  $w|_I$ ). Then, the free algebra  $k\langle\mathbb{A}\rangle$  is a Hopf algebra with comultiplication [31, 24].

$$\Delta_{LieHopf}(w) = \sum_{I+J=[1..n]} w[I] \otimes w[J]. \quad (14)$$

One has the following relation between restrictions of diagrams and subwords

$$\varphi_{lm}(d[I]) = cpt(\varphi_{lm}(d)[I]) \quad (15)$$

this suggests that the coproduct

$$\Delta_{list}(l) = \sum_{I+J=[1..n]} cpt(l[I]) \otimes cpt(l[J]) \quad (16)$$

could be a Hopf algebra comultiplication for the shifted algebra  $(k\langle\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X})\rangle, \bar{*}, [ ])$ . Unfortunately, this fails due to the lack of counit (i and ii of the following Theorem), but the “ground subalgebra”  $\mathcal{C}_{ldiag}$  is a genuine Hopf algebra (which is exactly what we do need here).

**Theorem 4.3** *Let  $\mathcal{A} = (k\langle \mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X}) \rangle, \bar{*}, [ \ ])$  be the algebra of lists of (non-unit) monomials endowed with the shifted concatenation of formula (10). Then*

*i)  $\mathcal{A}$  is a free algebra.*

*ii) The coproduct  $\Delta_{list}$  (recalled below) is co-associative and a morphism of algebras  $\mathcal{A} \mapsto \mathcal{A} \otimes \mathcal{A}$  (i.e.  $\mathcal{A}$  is a bi-algebra without counit).*

$$\Delta_{list}(l) = \sum_{I+J=[1..n]} cpt(l[I]) \otimes cpt(l[J]) \quad (17)$$

*iii) The algebra  $\mathcal{C}_{ldiag}$  is a sub- algebra and coalgebra of  $\mathcal{A}$  which is a Hopf algebra for the following co-unit and antipode.*

• COUNIT

$$\varepsilon(l) = \delta_{l, [ \ ]} \quad (\text{Kronecker delta}) \quad (18)$$

• ANTIPODE

$$S(l) = \sum_{r \geq 0} \sum_{\substack{I_1+I_2+\dots+I_r=[1..p] \\ I_j \neq \emptyset}} (-1)^r cpt(l[I_1])cpt(l[I_2]) \cdots cpt(l[I_r]) \quad (19)$$

*Proof* — i) Throughout the proof, we will denote  $*$  the concatenation between lists and  $\bar{*}$  the shifted concatenation defined by formula (10). Let's first remark that, if  $l = l_1 \bar{*} l_2$ , then  $\max(\text{IndAlph}(l_1)) < \min(\text{IndAlph}(l_2))$ . This leads us to define, for a (non-shifted) factorization  $l = l_1 * l_2 = l[1..t] * l[t+1..p]$  ( $p = |l|$ ) a gauge of the degree of overlapping of the intervals (of integers)  $[1..\max(\text{IndAlph}(l_1))]$  and  $[\min(\text{IndAlph}(l_2))..\infty[$ , thus the function

$$\omega_l(t) = \text{card} \left( [1..\max(\text{IndAlph}(l[1..t]))] \cap [\min(\text{IndAlph}(l[t+1..p]))..\infty[ \right) = \left( \max(\text{IndAlph}(l[1..t])) - \min(\text{IndAlph}(l[t+1..p])) + 1 \right)^+ \quad (20)$$

(we recall that, for a real number  $x$ ,  $x^+$  is its positive part i.e.  $x^+ = \max(x, 0) = \frac{1}{2}(|x| + x)$  [8]). It can be easily checked that the points  $t$  where  $\omega_l(t) = 0$  determine the (then unique) factorisation of  $l$  in irreducibles. It follows that the monoid  $((\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X}))^*, \bar{*}, [ \ ])$  is free and so is its algebra  $(k\langle \mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X}) \rangle, \bar{*}, [ \ ])$ .

ii) If we denote  $\Delta : \mathcal{A} \mapsto \mathcal{A} \otimes \mathcal{A}$  the standard coproduct given, for a list  $l$  of length  $p$ , by formula (14), one can remark that

- (i)  $cpt(l_1) \bar{*} cpt(l_2) = cpt(l_1 \bar{*} l_2)$
- (ii)  $\Delta_{list} = (cpt \otimes cpt) \circ \Delta$
- (iii)  $\Delta_{list} \circ cpt = \Delta_{list}$
- (iv)  $(\forall n \in \mathbb{N})(cpt(T_n(l)) = cpt(l))$
- (v)  $(\forall n \in \mathbb{N})(\Delta \circ T_n = (T_n \otimes T_n) \circ \Delta)$

COASSOCIATIVITY OF  $\Delta_{list}$ . —

One has

$$\begin{aligned}
 (\Delta_{list} \otimes Id) \circ \Delta_{list} &= (\Delta_{list} \otimes Id) \circ (cpt \otimes cpt) \circ \Delta = ' \\
 ((\Delta_{list} \circ cpt) \otimes cpt) \circ \Delta &= (\Delta_{list} \otimes cpt) \circ \Delta = \\
 (((cpt \otimes cpt) \circ \Delta) \otimes cpt) \circ \Delta &= \\
 (cpt \otimes cpt \otimes cpt) \circ (\Delta \otimes Id) \circ \Delta &= (cpt \otimes cpt \otimes cpt) \circ (Id \otimes \Delta) \circ \Delta \\
 (cpt \otimes ((cpt \otimes cpt) \circ \Delta)) \circ \Delta &= (cpt \otimes \Delta_{list}) \circ \Delta = \\
 (cpt \otimes (\Delta_{list} \circ cpt)) \circ \Delta &= \\
 (Id \otimes \Delta_{list}) \circ (cpt \otimes cpt) \circ \Delta &= (Id \otimes \Delta_{list}) \circ \Delta_{list}
 \end{aligned} \tag{21}$$

$\Delta_{list}$  IS A MORPHISM. —

For two lists  $u, v \in$ , let us compute  $\Delta_{list}(u\bar{*}v)$ . With  $p = \max(IndAlph(u))$ , one has

$$\begin{aligned}
 \Delta_{list}(u\bar{*}v) &= (cpt \otimes cpt) \circ \Delta(l_1 * T_p(v)) = \\
 (cpt \otimes cpt)(\Delta(u) *^{\otimes 2} \Delta(T_p(v))) &= \\
 (cpt \otimes cpt)(\Delta(u) *^{\otimes 2} (T_p \otimes T_p)\Delta(v)) &= \\
 (cpt \otimes cpt)\left(\sum_{(1)(2)} u_{(1)} \otimes u_{(2)} *^{\otimes 2} (T_p \otimes T_p)\left(\sum_{(3)(4)} v_{(3)} \otimes v_{(4)}\right)\right) &= \\
 (cpt \otimes cpt)\left(\sum_{(1)(2)(3)(4)} u_{(1)} * T_{p_1}(T_{p-p_1}(v_{(3)})) \otimes u_{(2)} * T_{p_2}(T_{p-p_2}(v_{(4)}))\right) & \tag{22}
 \end{aligned}$$

with, for each term in the sum

$$p_1 = \max(IndAlph(u_{(1)})) \leq p ; p_2 = \max(IndAlph(u_{(2)})) \leq p$$

so, the quantity in (22) is

$$\begin{aligned}
 (cpt \otimes cpt)\left(\sum_{(1)(2)(3)(4)} u_{(1)}\bar{*}(T_{p-p_1}(v_{(3)})) \otimes u_{(2)}\bar{*}(T_{p-p_2}(v_{(4)}))\right) &= \\
 \sum_{(1)(2)(3)(4)} cpt(u_{(1)}\bar{*}(T_{p-p_1}(v_{(3)}))) \otimes cpt(u_{(2)}\bar{*}(T_{p-p_2}(v_{(4)}))) &= \\
 \sum_{(1)(2)(3)(4)} \left(cpt(u_{(1)})\bar{*}cpt(T_{p-p_1}(v_{(3)}))\right) \otimes \left(cpt(u_{(2)})\bar{*}cpt(T_{p-p_2}(v_{(4)}))\right) &= \\
 \sum_{(1)(2)(3)(4)} \left(cpt(u_{(1)})\bar{*}cpt(v_{(3)})\right) \otimes \left(cpt(u_{(2)})\bar{*}cpt(v_{(4)})\right) &= \\
 \left(\sum_{(1)(2)} cpt(u_{(1)}) \otimes cpt(u_{(2)})\right)\bar{*}^{\otimes 2} \left(\sum_{(3)(4)} cpt(v_{(3)}) \otimes cpt(v_{(4)})\right) &= \\
 \Delta_{list}(u)\bar{*}^{\otimes 2} \Delta_{list}(v) & \tag{23}
 \end{aligned}$$

iii) As  $\mathcal{C}_{ldiag}$  is generated by the image of  $cpt$  it is clear that this space is a subcoalgebra of  $\mathcal{A}$ . Moreover,  $cpt$  is a (multiplicative) morphism  $\mathcal{A} \mapsto \mathcal{A}$  and thus its image

$\mathcal{C}_{ldiag}$  is a subalgebra of  $\mathcal{A}$ . Let us prove what is missing to make complete the Hopf algebra structure.

$\varepsilon$  IS A COUNIT. —

Let  $l = cpt(l)$  be a compact list. We remark that, for any list  $u$ , one has  $cpt(u) = [ ] \iff u = [ ]$ . Then, with  $\mu_l : k \otimes \mathcal{A} \mapsto \mathcal{A}$  the scaling operator

$$\begin{aligned} \mu_l(\varepsilon \otimes Id)\Delta_{list}(l) &= \sum_{I+J=[1..n]} \varepsilon(cpt(l[I]))cpt(l[J]) = \\ & \sum_{\substack{I+J=[1..n] \\ I=\emptyset}} \varepsilon(cpt(l[I]))cpt(l[J]) + \sum_{\substack{I+J=[1..n] \\ I \neq \emptyset}} \varepsilon(cpt(l[I]))cpt(l[J]) = cpt(l) + 0 = l \end{aligned} \quad (24)$$

the proof of the fact that  $\varepsilon$  is a left counit is similar.

$S$  IS THE ANTIPODE. —

One has  $\mathcal{C}_{ldiag} = k.1 \oplus ker(\varepsilon)$ , let us denote  $Id^+$  the projection  $\mathcal{C}_{ldiag} \mapsto ker(\varepsilon)$  according to this decomposition.

Then, for every list  $l$ ,

$$\sum_{r \geq 0} \sum_{\substack{I_1+I_2+\dots+I_r=[1..p] \\ I_j \neq \emptyset}} (-1)^r cpt(l[I_1])cpt(l[I_2]) \cdots cpt(l[I_r])$$

is well defined as the first sum is locally finite. Thus, the operator

$$\sum_{r \geq 0} \sum_{\substack{I_1+I_2+\dots+I_r=[1..p] \\ I_j \neq \emptyset}} (-1)^r \underbrace{(Id^+ * Id^+ * \cdots * Id^+)}_{r \text{ times}}$$

is well defined and is the convolutional inverse of  $Id$ .

### 4.3. Subalgebras of **LDIAG**

**4.3.1. Graphic primitive elements** The problem of Graphic Primitive Elements (GPE) is the following.

Let  $\mathcal{H}$  be a Hopf algebra with (linear) basis  $G$ , a set of graphs. The GPE are the primitive elements  $\Gamma \in G$  which are primitive i.e.

$$\Gamma \text{ is a GPE} \iff \Gamma \in G \text{ and } \Delta(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma \quad (25)$$

it is not difficult to check that, in any case, the subalgebra  $\mathcal{H}^{GPE}$  generated by these elements is also a sub-coalgebra.

We do the extra hypothesis (which is often fulfilled)

$$1_{\mathcal{H}} \in G \text{ and } (\Gamma \in G - \{1_{\mathcal{H}}\} \implies \varepsilon(\Gamma) = 0). \quad (26)$$


Then (if (26) is fulfilled)  $\mathcal{H}^{GPE}$  is a sub-Hopf algebra as the antipode of a product  $\Gamma_1 \Gamma_2 \cdots \Gamma_p$  of (GPE) is

$$S(\Gamma_1 \Gamma_2 \cdots \Gamma_p) = (-1)^p \Gamma_p \Gamma_{p-1} \cdots \Gamma_1. \quad (27)$$

The following proposition helps to determine **LDIAG**<sup>GPE</sup>.

**Proposition 4.4** *In **LDIAG** (with basis  $G = \mathbf{ldiag}$ ), the following are equivalent*  
*i)  $d$  is a GPE*  
*ii)  $d$  has only one black spot.*

Then, the Hopf algebra  $\mathbf{LDIAG}^{\text{GPE}}$  is generated by the product of “one-black-spot” diagrams.



**Fig 4.** — *Graphic Primitive Elements of **LDIAG** have only one black spot and therefore are coded by the sequence of the ingoing degrees of their white spots (a composition). The first one here has code  $[1, 2, 3, 1]$ . The picture shows an element of the monoid generated by Graphic Primitive Elements (a linear basis of  $\mathbf{LDIAG}^{\text{GPE}}$ ) which is then coded by a list of compositions, here  $\left[ [1, 2, 3, 1], [2, 3, 1], [2, 1, 4] \right]$ .*

#### 4.3.2. Level subalgebras

One can impose also limitations on the ingoing degrees of the white spots in a way compatible with the coproduct. In this case, one defines an infinity of Hopf-subalgebras of **LDIAG** which we will call “level subalgebras”.

More precisely, given an integer  $l > 0$ , one can ask for spaces generated by the diagrams  $d$  for which every white spot has an ingoing degree  $\leq l$ . This amounts to say that the “white spots type” of every diagram  $d$  is of the form

$$\alpha(d) = (\alpha_1, \alpha_2, \dots, \alpha_k, 0, 0 \dots 0, \dots) ; \quad (\text{all the } \alpha_i \leq l \text{ for } i \leq k \text{ and } \alpha_i = 0 \text{ for } i > k)$$

Let us denote  $\mathbf{LDIAG}^{\leq l}$  the subspace generated by these diagrams. One has a chain of Hopf algebras

$$\mathbf{LDIAG}^{\leq 1} \subset \mathbf{LDIAG}^{\leq 2} \subset \dots \mathbf{LDIAG}^{\leq l} \subset \mathbf{LDIAG}^{\leq l+1} \subset \dots \subset \mathbf{LDIAG} \quad (28)$$

In the next paragraph, we will specially be interested by

$$\mathbf{LBELL} = \mathbf{LDIAG}^{\leq 1} \cap \mathbf{LDIAG}^{\text{GPE}}.$$

#### 4.3.3. BELL and LBELL


The algebras **BELL** and **LBELL** were defined in [36].

be done : following Allan, explain the link with coherent states.

The algebra **LBELL** is the intersection  $\mathbf{LDIAG}^{\leq 1} \cap \mathbf{LDIAG}^{\text{GPE}}$  and then, as they are subspaces generated by subsets of **ldiag**, **LBELL** is generated by diagrams that

- are concatenation of one-black-spot-diagrams
- the ingoing degree of every white spot is one.

Let  $d_k$  be the diagram with code  $[x_1, x_2, \dots, x_k]$ . **LBELL** is generated by concatenations of these diagrams. Indeed, the diagrams  $d_k$  are a subalphabet of the free monoid **ldiag** so that they generate a free submonoid which we will denote here **lbell**.



**Fig 5.** — An element of **lbell**, concatenation  $d_1d_3d_2$ .

The algebras **LDIAG** and **LBELL** are both envelopping algebras. They generated by their primitive elements which are in general linear combination of diagrams and not pure diagrams. For an analysis of “graphic primitive elements” see section (4.3.1).

## 5. The algebra **LDIAG**( $q_c, q_s, q_t$ ) (deformed case)

### 5.1. Counting crossings ( $q_c$ ) and superpositions ( $q_s$ )


The preceding coding is particularly well adapted to the the deformation we want to construct here. The philosophy of the deformed product is expressed by the descriptive formula.

$$[d_1|d_2]_{L(q_c, q_s)} = \sum_{\substack{cs(?) \text{ all crossing and} \\ \text{superpositions of black spots}}} q_c^{nc \times weight} q_s^{weight \times weight} cs([d_1|d_2]_L) \quad (29)$$


where

- $q_c, q_s \in \mathbb{C}$  or  $q_c, q_s$  formal. These and other cases being unified with considering the set of coefficients as taken in a ring  $K$ .
- the exponent of  $q_c^{nc \times weight}$  is the number of crossings of “what crosses” times its weight
- the exponent of  $q_s^{weight \times weight}$  is the product of the weights of “what is overlapped”
- $cs(?)$  are the diagrams obtained from  $[d_1|d_2]_L$  by the process of crossing and superposing the black spots of  $d_2$  to those of  $d_1$ , the order and distinguishability of the black spots of  $d_1$  (i.e.  $d_2$ ) being preserved.

What is striking is that this law is associative. This result will be established after the following paragraph.



**Fig 5.** — Counting crossings and superposings produces an associative law.



**Fig 6.** — Detail of the fourth monomial (with coefficient  $q_c^2 q_s^6$ ), crossings (circles) and superposings (black squares) are counted the same way but with a different variable.

5.2. Modified laws

Introduction : Dual laws, shuffle, infiltration, Hoffmann.

- Twisting

**Proposition 5.1** Let  $A = (A_n)_{n \in \mathbb{N}}$  a graded semigroup and  $A^*$  the set of lists (denoted  $[a_1, a_2, \dots, a_k]$ ) with letters in  $A$ .

For convenience we define the operator  $*$  (left append)  $A \times A^* \mapsto A^*$  by

$$a * [b_1, b_2, \dots, b_n] := [a, b_1, b_2, \dots, b_n] \tag{30}$$



Let  $q_c, q_s \in k$  be two elements in a ring  $k$ . We define on  $k \langle A \rangle = k[A^*]$  a new law  $\uparrow$  by

$$\begin{aligned} w \uparrow 1_{A^*} &= 1_{A^*} \uparrow w = w \\ a * u \uparrow b * v &= a * (u \uparrow b * v) + q_c^{|a*u||b|} b * (a * u \uparrow v) + q_c^{|u||b|} q_s^{|a||b|} ab * (u \uparrow v) \end{aligned} \quad (31)$$

where the weights ( $|x| = n$  if  $x \in A_n$ ) are extended additively to lists by

$$|[a_1, a_2, \dots, a_k]| = \sum_{i=1}^k |a_i|$$

Then the new law  $\uparrow$  is graded, associative with  $1_{A^*}$  as unit.

*Proof* — It suffices to prove the identity  $x \uparrow (y \uparrow z) = (x \uparrow y) \uparrow z$ ;  $x, y, z$  being lists (as the two members are trilinear). It is obviously true when one of the factors is the void list. Let's show it when the three factors are non-void (throughout the computation, the law  $*$  will have priority over other operators).

$$\begin{aligned} &(a * u \uparrow b * v) \uparrow c * w = \\ &(a * (u \uparrow b * v) + q^{|u||b|} t^{|a||b|} (ab)(u \uparrow v) + q^{|a*u||b|} b(a * u \uparrow v)) \uparrow c * w = \\ &\left[ a * ((u \uparrow b * v) \uparrow c * w) + q^{(|u|+|b*v|)|c|} t^{|a||c|} (ac)((u \uparrow b * v) \uparrow w) \right. \\ &\left. + q^{(|a*u|+|b*v|)|c|} c * (a * (u \uparrow b * v) \uparrow w) \right] + \\ &\left[ q^{|u||b|} t^{|a||b|} (ab)(u \uparrow v \uparrow c * w) + q^{|u||b|+(|u|+|v|)|c|} t^{|a||b|} t^{(|a|+|b|)|c|} (abc)(u \uparrow v \uparrow w) + \right. \\ &\left. q^{|u||b|+(|a*u|+|b*v|)|c|} t^{|a||b|} c(((ab)(u \uparrow v)) \uparrow w) \right] + \\ &\left[ q^{|a*u||b|} b((a * u \uparrow v) \uparrow c * w) + q^{|a*u||b|+(|a*u|+|v|)|c|} t^{|b||c|} (bc)(au \uparrow v \uparrow w) + \right. \\ &\left. q^{|a*u||b|+(|a*u|+|b*v|)|c|} c(b(a * u \uparrow v) \uparrow w) \right] \end{aligned} \quad (32)$$

$$\begin{aligned} &a * u \uparrow (b * v \uparrow c * w) = \\ &a * u \uparrow (b * (v \uparrow c * w) + q^{|v||c|} t^{|b||c|} (bc)(v \uparrow w) + q^{|b*v||c|} c(b * v \uparrow w)) = \\ &\left[ a * (u \uparrow b * (v \uparrow c * w)) + q^{|u||b|} t^{|a||b|} (ab)(u \uparrow v \uparrow c * w) + q^{|a*u||b|} b(a * u \uparrow v \uparrow c * w) \right] + \\ &\left[ q^{|v||c|} t^{|b||c|} a * (u \uparrow (bc)(v \uparrow w)) + q^{|v||c|+|u|(|c|+|b|)} t^{|b||c|+|a|(|b|+|c|)} (abc)(u \uparrow v \uparrow w) + \right. \\ &\left. q^{|v||c|+|a*u|(|b|+|c|)} t^{|b||c|} (bc)(a * u \uparrow v \uparrow w) \right] + \\ &\left[ q^{|b*v||c|} a * (u \uparrow c(b * v \uparrow w)) + q^{(|u|+|b*v|)|c|} t^{|a||c|} (ac)(u \uparrow b * v \uparrow w) + \right. \\ &\left. q^{(|a*u|+|b*v|)|c|} c * (a * u \uparrow b * v \uparrow w) \right] \end{aligned} \quad (33)$$

in the second expression, one gathers the three terms which we find first in the square brackets and we get

$$a * (u \uparrow b * (v \uparrow cw)) + q^{|v||c|} t^{|b||c|} a * (u \uparrow (bc) * (v \uparrow w)) + q^{|b*v||c|} a * (u \uparrow c * (b * v \uparrow w)) = a * (u \uparrow b * v \uparrow c * w) \quad (34)$$

in the first expression, one gathers the three terms which we find last in the square brackets and we get

$$\begin{aligned} & q^{(|a*u|+|b*v|)|c|} c * (a * (u \uparrow b * v) \uparrow w) + \\ & q^{|u||b|+(|a*u|+|b*v|)|c|} t^{|a||b|} c * (((ab) * (u \uparrow v)) \uparrow w) + \\ & q^{|a*u||b|+(|a*u|+|b*v|)|c|} c * (b * (a * u \uparrow v) \uparrow w) = \\ & q^{(|au|+|bv|)|c|} c * (a * u \uparrow b * v \uparrow w) \end{aligned} \quad (35)$$

and one finds the 7 terms expression

$$\begin{aligned} & a * (u \uparrow b * v \uparrow c * w) + q^{|a*u|} b * (a * u \uparrow v \uparrow c * w) + \\ & q^{|a*u|+|b*v|} c * (a * u \uparrow b * v \uparrow w) + q^{|u||b|} t^{|a||b|} (ab) * (u \uparrow v \uparrow c * w) + \\ & q^{(|u|+|b*v|)|c|} t^{|a||c|} (ac) * (u \uparrow b * v \uparrow w) + \\ & q^{|v||c|(|b|+|c|)|au|} t^{|b||c|} (bc) * (a * u \uparrow v \uparrow w) \\ & + q^{|v||c|+|u|(|c|+|b|)} t^{|b||c|+|a|(|b|+|c|)} (abc) * (u \uparrow v \uparrow w) \end{aligned} \quad (36)$$

□

The framework with diagrams will need another proposition on shifted laws.

- *Shifting*

We begin by the “shifting lemma” [19].

**Lemma 5.2** *Let  $\mathcal{A}$  be an associative algebra (which law will be denoted  $\star$ ) and  $\mathcal{A} = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n$  a decomposition of  $\mathcal{A}$  in direct sum. Let  $T \in \text{End}(\mathcal{A})$  be an endomorphism of the algebra  $\mathcal{A}$ . We will denote  $T^n = T \circ T \circ \dots \circ T$  the  $n$ -th compositional power of  $T$ . We suppose that the shifted law*

$$a \bar{\star} b = a \star T^\alpha(b) \quad (37)$$

for  $a \in \mathcal{A}_\alpha$  is graded for the decomposition  $\mathcal{A} = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n$ .

Then, if the law  $\star$  is associative so is the law  $\bar{\star}$ .

**Remark 5.3** *The hypothesis that the shifted law given by eq.(37) be graded is automatically satisfied if  $\mathcal{A} = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n$  is a graded algebra and if all the morphisms  $T_n$  are of degree 0.*

This lemma will be applied to the decomposition given by  $n = \text{sup}(\text{Alph}(w))$  (the highest index of variables appearing in  $w$ ) and the morphism given by  $T(x_i) = x_{i+1}$ .

What does these statements mean for us ?

Here the graded semigroup is  $\mathfrak{MON}^+(X)$  and we do not forget the coding arrow

$\varphi_{lm} : \mathbf{ldiag} \rightarrow (\mathfrak{M}\mathfrak{D}\mathfrak{N}^+(X))^*$ . The image of  $\varphi_{lm}$  is exactly the set of lists of monomials  $w = [m_1, m_2, \dots, m_k]$  such that the set of variables involved  $Alph(w)$  is of the form  $x_1 \cdots x_l$  (the labelling of the white spots is without hole). By abuse of language we will say that a list of monomials “is in  $\mathbf{ldiag}$ ” in this case. It is not difficult to see, from formulas (31,37) that if  $w_i$ ,  $i = 1, 2$  are in  $\mathbf{ldiag}$  so are all the factors of  $w_1 \bar{\uparrow} w_2$ , this defines a new law on  $K[\mathbf{ldiag}]$  and this algebra will be called  $\mathbf{LDIAG}(q_c, q_s)$ . The properties of this algebra will be made precise in the following proposition.

**Proposition 5.4** *Let  $\mathcal{C}_{ldiag}$  be the subspace of  $(K < \mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X}) >, \bar{\uparrow})$  generated by the codes of the diagrams (i.e. the lists  $w \in \mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X})$  such that  $Alph(w)$  is without hole).*

*Then*

- i)  $(\mathcal{C}_{ldiag}, \bar{\uparrow})$  is a unital subalgebra of  $(K < \mathfrak{M}\mathfrak{D}\mathfrak{N}^+(\mathbb{X}) >, \bar{\uparrow})$*
- ii)  $(\mathcal{C}_{ldiag}, \bar{\uparrow})$  is a free algebra. More precisely, for any diagram decomposed in irreducibles  $d = d_1.d_2 \cdots d_k$  let*

$$B(d) := \varphi_{lm}(d_1) \bar{\uparrow} \varphi_{lm}(d_2) \cdots \bar{\uparrow} \varphi_{lm}(d_k) \tag{38}$$

*then*

- $\alpha)$   $(B(d))_{d \in ldiag}$  is a basis of  $\mathcal{C}_{ldiag}$
- $\beta)$   $B(d_1.d_2) = B(d_1) \bar{\uparrow} B(d_2)$

As  $k[\mathbf{ldiag}]$  is isomorphic to  $\mathcal{C}_{ldiag}$  as a linear space, we denote  $\mathbf{LDIAG}(q_c, q_s)$  the new algebra structure of  $k[\mathbf{ldiag}]$  inherited from  $\mathcal{C}_{ldiag}$ . one has

$$\mathbf{LDIAG}(0, 0) \simeq \mathbf{LDIAG}; \quad \mathbf{LDIAG}(1, 1) \simeq \mathbf{MQSym} \tag{39}$$

## 6. Coproducts

Now, one has to define a parametrized (by, say,  $q_t$ ) coproduct such that  $(\mathbf{LDIAG}(q_c, q_s), \bar{\uparrow}, \mathbf{1}_{\mathbf{ldiag}}, \Delta_{q_t}, \varepsilon)$  be a graded bialgebra (the counity  $\varepsilon$ , the same as in the non-deformed Hopf algebra in [14] is just the “constant term” linear form). We will take advantage of the freeness of  $\mathbf{LDIAG}(q_c, q_s)$  through the following lemma.

**Lemma 6.1** *Let  $\mathbb{Y}$  be an alphabet,  $k$  a ring and*

*$k < \mathbb{Y} > = k[\mathbb{Y}^*]$  be the free algebra constructed on  $\mathbb{Y}$ . For every mapping*

*$\Delta : A \rightarrow k < \mathbb{Y} > \otimes k < \mathbb{Y} >$ , we denote  $\bar{\Delta} : k < \mathbb{Y} > \mapsto k < \mathbb{Y} > \otimes k < \mathbb{Y} >$  its extension as a morphism of algebras ( $k < \mathbb{Y} > \otimes k < \mathbb{Y} >$  being endowed with its non-twisted structure of tensor product of algebras). Then, in order be coassociative, it is necessary and sufficient that*

$$(\bar{\Delta} \otimes I) \circ \Delta \text{ and } (I \otimes \bar{\Delta}) \circ \Delta \tag{40}$$

*coincide on  $\mathbb{Y}$ .*

The preceding lemma expresses that, for a free algebra, the “phase space” of the possible coproducts is a linear subspace. This will be transparent in formula (43).

Now, we consider the structure constants of the coproduct of **MQSym** [17] expressed with respect to the family of free generators

$$\{MS_P\}_{P \in \mathcal{PM}^c}$$

where  $\mathcal{PM}^c$  is the set of connex packed matrices (similarly,  $\mathcal{PM}$  is the set of packed matrices).

$$\Delta_{\mathbf{MQSym}}(MS_P) = \sum_{Q,R \in \mathcal{PM}} \alpha_P^{Q,R} MS_Q \otimes MS_R \quad (41)$$

For  $d$ , irreducible diagram, set

$$\Delta_1(d) = \sum_{d_1, d_2 \in \text{irr}(\mathbf{ldiag})} \alpha_{\varphi_{lm}(d)}^{\varphi_{lm}(d_1), \varphi_{lm}(d_2)} d_1 \otimes d_2 \quad (42)$$

and  $\Delta_0(d) = \Delta_{WS}(d)$ . Then proposition (6.1) proves that

$$\Delta_t = \overline{(1 - q_t)\Delta_0 + q_t\Delta_1} \quad (43)$$

is a coproduct of graded bialgebra for  $(\mathbf{LDIAG}(q_c, q_s), \uparrow, \mathbf{1}_{\mathbf{ldiag}})$ .

We sum up the results

**Proposition 6.2** *i) With the operations defined above*

$$\mathbf{LDIAG}(q_c, q_s, q_t) := (\mathbf{LDIAG}(q_c, q_s), \uparrow, \mathbf{1}_{\mathbf{ldiag}}, \Delta_{q_t}, \varepsilon)$$

*is a Hopf algebra.*

*ii) At parameters  $(0, 0, 0)$ , one has  $\mathbf{LDIAG}(0, 0, 0) \simeq \mathbf{LDIAG}$*

*iii) At parameters  $(1, 1, 1)$ , one has  $\mathbf{LDIAG}(1, 1, 1) \simeq \mathbf{MQSym}$*

**Comment 6.3** (The following comment could appear as ‘‘concluding remarks’’ what do you think ?)

*For a diagram  $d$  with  $r$  black spots, the code  $[m_1, m_2, \dots, m_r]$  can be temporarily seen as a ‘‘vector of coordinates’’ for the given diagram, but we preferred to stick to the structure of list as, firstly, the dimension of the vector varies with the diagram and secondly, we had to concatenate the codes. The coordinate functions of the diagram  $d$  are therefore the family  $(a_i)_{i \geq 0}$  defined by  $a_i(d) = m_i$  for  $i \leq r$  and  $a_i(d) = 0$  for  $i > r$ . In this perspective the ‘‘ $q_t$ ’’ of our three parameters deformation is a quantization in the sense of Moyal’s deformed products [1] on the algebra of coordinate functions (but without first order condition, for this see also the introduction of [12]), by the formula*

$$a_{i_1} * a_{i_1} \cdots * a_{i_k}(d) = \mu(a_{i_1} \otimes a_{i_1} \otimes \cdots \otimes a_{i_k}(\Delta_{q_t}^{[k]}(d))) \quad (44)$$

*where  $\mu$  is the ordinary multiplication of polynomials.*

*The crossing parameter  $q_c$  is also a quantization parameter as, for  $q_s = 0$ , one has*

$$\text{code}(d_1 * d_2) = \text{code}(d_1) \sqcup_{q_c} T(\text{code}(d_2)) \quad (45)$$

where  $T$  is a suitable translation of the variables and  $\sqcup_{q_c}$  is the quantum shuffle [32] for the braiding on  $V = \mathbb{C}[x_i; i \geq 1]$  defined by

$$B(x_{i_1}^{\alpha_1} x_{i_2}^{\alpha_2} \cdots x_{i_k}^{\alpha_k} \otimes y_{j_1}^{\beta_1} y_{j_2}^{\beta_2} \cdots y_{j_l}^{\beta_l}) = q_c^{(\sum \alpha_i)(\sum \beta_j)} y_{j_1}^{\beta_1} y_{j_2}^{\beta_2} \cdots y_{j_l}^{\beta_l} \otimes x_{i_1}^{\alpha_1} x_{i_2}^{\alpha_2} \cdots x_{i_k}^{\alpha_k} \quad (46)$$

Let us add that  $q_s$  and  $q_c$  are structurally different as  $q_s$  is the coefficient of a perturbation of the shuffle product (better seen on the coproduct). This kind of perturbation occurs in various domains as : computer science by means of the infiltration product introduced by Ochsenschläger [30] (see also [16] and [15]), algebra of the Euler-Zagier sums [25] and noncommutative symmetric functions [17]. The mathematics of this dual aspect is of geometrical nature and will be developed in [18].

## 7. LDIAG( $q_c, q_s, q_t$ ) and its specializations

Graphic primitive elements, BELL, LBELL, Zagier, Foissy and the general picture.

## 8. Conclusion

Discuss the self-dual deformation.

## References

- [1] BAYEN (F.), FLATO (M.), FRONSDAL (C.), LICHNEROWICZ (A.) AND STERNHEIMER (D.), *Deformation and Quantization*, Ann. of Phys. 111 (1978), pp. 61-151.
- [2] J. BERSTEL, C. REUTENAUER, *Rational series and their languages EATCS Monographs on Theoretical Computer Science*, Springer (1988).
- [3] C. M. BENDER, D. C. BRODY, AND B. K. MEISTER, Quantum field theory of partitions, J. Math. Phys. Vol 40 (1999)
- [4] C. BERTELLE, G. H. E. DUCHAMP, K. KHATATNEH, *Tables, Memorized Semirings and Applications*, arXiv: cs.MA/0502081.
- [5] P. BLASIAK, A. HORZELA, K. A. PENSON, G. H. E. DUCHAMP, A.I. SOLOMON, *Boson normal ordering via substitutions and Sheffer-Type Polynomials*, Phys. Lett. A **338** (2005) 108
- [6] P. BLASIAK, K. A. PENSON, A.I. SOLOMON, A. HORZELA, G. H. E. DUCHAMP, *Some useful formula for bosonic operators*, Jour. Math. Phys. **46** 052110 (2005).
- [7] BOURBAKI N., *Algebra, chapter III*, Springer
- [8] BOURBAKI N., *Algebra, chapter VI*, Springer
- [9] BOURBAKI N., *Theory of sets*, Springer
- [10] CARTIER P., *Séminaire "Sophus Lie", 2ème année*, Faculté des Sciences de Paris (1955-56)
- [11] P. CARTIER, *Fonctions polylogarithmes, nombres polyzeta et groupes pro-unipotents*, *Séminaire Bourbaki, Asterisque n. 282* (2002)
- [12] V. CHARI, A. PRESSLEY, *A guide to quantum groups*. Cambridge (1994).
- [13] L. COMTET, *Analyse Combinatoire*, PUF, Tome 1-2 (1970)
- [14] G. H. E. DUCHAMP, P. BLASIAK, A. HORZELA, K. A. PENSON, A. I. SOLOMON, FEYNMAN GRAPHS AND RELATED HOPF ALGEBRAS, *Journal of Physics: Conference Series*, SSPCM'05, Myczkowce, Poland. arXiv : cs.SC/0510041
- [15] DUCHAMP G., FLOURET M., LAUGEROTTE E., LUQUE J-G., *Direct and dual laws for automata with multiplicities*  
arXiv : math.CO0607412

- [16] DUCHAMP G., LUQUE J-G., *Congruences Compatible with the Shuffle Product*  
arXiv : math.CO0607419
- [17] G. DUCHAMP, F. HIVERT, J. Y. THIBON, *Non commutative functions VI: Free quasi-symmetric functions and related algebras*, International Journal of Algebra and Computation Vol 12, No 5 (2002).
- [18] G. H. E. DUCHAMP, G. KOSHEVOY, K. A. PENSON, C. TOLLU, F. TOUMAZET, *Geometric combinatorial twisting and shifting*, Séminaire Lotharingien (in preparation).
- [19] G. H. E. DUCHAMP, J. -G. LUQUE, J. -C. NOVELLI, C. TOLLU, F. TOUMAZET, *Hopf algebras of diagrams*, submitted to FPSAC07.
- [20] G. DUCHAMP, A.I. SOLOMON, K.A. PENSON, A. HORZELA AND P. B LASIAK, *One-parameter groups and combinatorial physics*, Proceedings of the Symposium Third International Workshop on Contemporary Problems in Mathematical Physics (COPROMAPH3) (Porto-Novo, Benin, Nov. 2003), J. Govaerts, M. N. Hounkonnou and A. Z. Msezane (eds.), p.436 (World Scientific Publishing 2004)  
arXiv: quant-ph/04011262
- [21] P. FLAJOLET, <http://algo.inria.fr/flajolet/>
- [22] L. FOISSY, *Isomorphisme entre l'algèbre des fonctions quasi-symétriques libres et une algèbre de Hopf des arbres enracinés décorés plans*, personal communication.
- [23] L. FOISSY, *Les algèbres de Hopf des arbres enracinés décorés*, PhD Memoir, Reims University (2002).
- [24] M. HAZEWINKEL, *Hopf algebras of endomorphisms of Hopf algebras*, (Oct 2004) ArXiv : math.QA/0410364
- [25] M. E. HOFFMAN, *Quasi-shuffle products*, J. Algebraic Combin. (2000), 49-68
- [26] A. HORZELA, P. BLASIAK, G. DUCHAMP, K. A. PENSON AND A.I. SOLOMON, *A product formula and combinatorial field theory*, Proceedings of the XI International Conference on Symmetry Methods in Physics (SYMPHYS-11) (Prague, Czech Republic, June 2004), C. Burdik, O. Navratil, and S. Posta (eds.) (JINR Publishers, Dubna)  
arXiv:quant-ph/0409152
- [27] S. A. JONI, G.-C.ROTA, *Colgebras and Bialgebras in Combinatorics*, Studies in Applied Mathematics **61**, 93-139 (1979).
- [28] D. KNUTH, *The art of computer programming* Tome I. Addison-Wesley (1981)
- [29] J. MILNOR, J. MOORE, *On the structure of Hopf Algebras*, Ann. of Math. 81 (1965), 211-264.
- [30] P. OCHSENSCHLÄGER, *Binomialkoeffizienten und Shuffle-Zahlen*, Technischer Bericht, Fachbereich Informatik, T. H. Darmstadt,1981.
- [31] REUTENAUER C., *Free Lie algebras*, Oxford University Press (1993).
- [32] M. ROSSO, *Quantum groups and quantum shuffles*, Inventiones Math. 133 (1998).
- [33] A.I. SOLOMON, P. BLASIAK, G. DUCHAMP, A. HORZELA AND K.A. PENSON, *Combinatorial Physics, Normal Order and Model Feynman Graphs*, Proceedings of the Symposium 'Symmetries in Science XIII', Bregenz, Austria, 2003, B. Gruber, G. Marmo and N. Yoshinaga (eds.), p.527 (Kluwer Academic Publishers 2004)  
arXiv : quant-ph/0310174
- [34] A.I. SOLOMON, G. DUCHAMP, P. BLASIAK, A. HORZELA AND K.A. PENSON, *Normal Order: Combinatorial Graphs Quantum Theory and Symmetries*, Proceedings of the 3rd International Symposium P.C. Argyres, T.J. Hodges, F. Mansouri, J.J. Scanio, P. Suranyi, and L.C.R. Wijewardhana (eds.), p.398 (World Scientific Publishing 2004)  
arXiv:quant-ph/0402082
- [35] A.I. SOLOMON, G. DUCHAMP, P. BLASIAK, A. HORZELA AND K. A. PENSON, *Partition functions and graphs: A combinatorial approach*, Proceedings of the XI International Conference on Symmetry Methods in Physics (SYMPHYS-11) (Prague, Czech Republic, June 2004), C. Burdik, O. Navratil, and S. Posta (eds.) (JINR Publishers, Dubna, 2004)  
arXiv:quant-ph/0409082

- [36] Allan's NY
- [37] Gerard's NY
- [38] R. P. STANLEY, *Enumerative Combinatorics*, vol. **2**, Cambridge Univ. Press (2004).
- [39] A. VARVAK, Rook numbers and the normal ordering problem, *Preprint* arXiv: **math.**  
**CO/0402376**
- [40] S. WEINZIERL, *Hopf algebra structures in particle physics*, Eur. Phys. J. C **33** (2004) arXiv:  
**hep-th/0310124**