

# Functional Stepped Surfaces, Flips and Generalized Substitutions

Pierre Arnoux,

*IML – CNRS UMR 6206– 163, avenue de Luminy, Case 907, 13288 Marseille  
Cedex 09, France*

Valérie Berthé, Thomas Fernique,

*LIRMM – UMR 5506 – Univ. Montpellier II, 161 rue Ada, 34392 Montpellier  
Cedex 05, France*

Damien Jamet,

*LORIA – INRIA Lorraine – Technopole de Nancy-Brabois, 615 rue du Jardin  
Botanique, 54600 Villers-lès-Nancy, France*

---

## Abstract

A substitution is a non-erasing morphism of the free monoid. The notion of multidimensional substitution of non-constant length acting on multidimensional words introduced in [AI01,ABS04,Fer05a,Fer05b,Fer05c] is proved to be well-defined on the set of two-dimensional words related to discrete approximations of irrational planes. Such a multidimensional substitution can be associated with any usual unimodular substitution. The aim of this paper is to extend the domain of definition of such multidimensional substitutions to functional stepped surfaces, such as defined in [Jam04,JP05]. One central tool for this extension is the notion of flips acting on tilings by lozenges of the plane.

*Key words:* Generalized substitution, discrete geometry, arithmetic discrete plane, discrete surface, word combinatorics, flips, lozenge tiling, dimer tiling, Sturmian word.

---

*Email addresses:* [arnoux@iml.univ-mrs.fr](mailto:arnoux@iml.univ-mrs.fr) (Pierre Arnoux), [berthe@lirmm.fr](mailto:berthe@lirmm.fr) (Valérie Berthé), [fernique@lirmm.fr](mailto:fernique@lirmm.fr) (Thomas Fernique), [jamet@loria.fr](mailto:jamet@loria.fr) (Damien Jamet).

## 1 Introduction

Sturmian words are known to be codings of digitizations of an irrational straight line [KR04,Loth02]. One could expect from a higher-dimensional generalization of Sturmian words that they correspond to a digitization of a hyperplane with irrational normal vector. It is thus natural to consider the digitization scheme corresponding to the notion of arithmetic planes introduced in [Rev91]: this notion consists in approximating a plane in  $\mathbb{R}^3$  by selecting points with integral coordinates above and within a bounded distance of the plane; more precisely, given  $\mathbf{v} \in \mathbb{R}^3$ , and  $(\mu, \omega) \in \mathbb{R}^2$ , the *lower* (resp. *upper*) *arithmetic hyperplane*  $\mathfrak{P}(\mathbf{v}, \mu, \omega)$  is the set of points  $\mathbf{x} \in \mathbb{Z}^3$  satisfying  $0 \leq \langle \mathbf{x}, \mathbf{v} \rangle + \mu < \omega$  (resp.  $0 < \langle \mathbf{x}, \mathbf{v} \rangle + \mu \leq \omega$ ). Moreover, if  $\omega = \sum |v_i| = \|\mathbf{v}\|_1$ , then  $\mathfrak{P}(\mathbf{v}, \mu, \omega)$  is said to be *standard*.

In this latter case, one approximates a plane with normal vector  $\mathbf{v} \in \mathbb{R}^3$  by square faces oriented along the three coordinates planes. The union of all these faces is called a *stepped plane*; the standard discrete plane  $\mathfrak{P}(\mathbf{v}, \mu, \|\mathbf{v}\|_1)$  is then equal to the set of points with integer coordinates that belong to the stepped plane; after orthogonal projection onto the plane  $x_1 + x_2 + x_3 = 0$ , one obtains a tiling of the plane with three kinds of lozenges, namely the projections of the three possible faces. One can code this projection over  $\mathbb{Z}^2$  by associating with each lozenge the name of the projected face it corresponds to. These words are in fact three-letter two-dimensional Sturmian words (see, e.g., [BV00]).

A generalization of the notion of stepped plane, the so-called *discrete surfaces*, is introduced in [Jam04,JP05]. A discrete surface is defined as a union of pointed faces such that the orthogonal projection onto the diagonal plane  $x_1 + x_2 + x_3 = 0$  induces an homeomorphism from the discrete surface onto the diagonal plane. As it is done for stepped planes, one provides any discrete surface with a coding as a two-dimensional word over a three-letter alphabet. In the present paper, we call the discrete surfaces *functional stepped surfaces*, since such objects are not discrete, in the sense that they are not subsets of  $\mathbb{Z}^3$ . Furthermore, one could define more general stepped surfaces, for instance approximations of spheres; the surfaces we consider here are functional, that is, they project homeomorphically onto the diagonal plane and can be described as graphs of piecewise affine maps defined on the diagonal plane.

Let us recall that a substitution is a non-erasing morphism of the free monoid. It acts naturally on all finite and infinite words. In particular, it maps a two-sided word to a two-sided word. We are interested here in higher dimensional analogues of substitutions. It is easy to define a two-dimensional substitution which replaces each letter by a rectangle of fixed size. This is the analogue of substitutions of constant length, and such a substitution acts on the set of

all two-dimensional words. For such examples, see for instance [AS03]. In the present paper, we deal with substitutions of non-constant length; one easily sees that such a substitution cannot be meaningfully defined on all the two-dimensional words: if two letters are replaced by patterns of different shapes, and if we consider two two-dimensional words that differ in exactly one place by the corresponding letters, it is not possible that both two-dimensional words are sent by the substitution to complete two-dimensional words. In fact, it is not even clear that the substitution can act on some two-dimensional words.

A notion of multidimensional substitution of non-constant length acting on multidimensional words is studied in [AI01,AIS01,ABI02,ABS04,Fer05b], inspired by the geometrical formalism of [IO93,IO94]. These multidimensional substitutions are proved in [AI01] to be well-defined on multidimensional Sturmian words. Such a multidimensional substitution can be associated with any usual unimodular substitution (a substitution is said *unimodular* if the determinant of its incidence matrix equals  $\pm 1$ ). The aim of the present paper is to explore the domain of definition of such multidimensional substitutions. The main result of the present paper is thus the following: the image of a functional stepped surface under the action of a 2-dimensional substitution is still a functional stepped surface.

Our proofs are based on a geometrical approach, using the generation of functional stepped surfaces by *flips*. A flip is a classical notion in the study of dimer tilings and lozenge tilings associated with the triangular lattice; see, e.g., [Thu89]. It consists in a local reorganization of tiles that transforms a tiling into another one. Such a reorganization can also be seen in the 3-dimensional space on the functional stepped surface itself. Suppose indeed that a functional stepped surface contains 3 faces that form the lower faces of a unit cube with integer vertices. By replacing these three faces by the upper faces of this cube, one obtains another functional stepped surface (see Figure 7). We prove that any functional stepped surface can be obtained from a stepped plane by a sequence of flips, possibly infinite but locally finite, in the sense that, for any bounded neighborhood of the origin in the diagonal plane, there is only a finite number of flips whose domain has a projection which intersects this neighborhood.

This paper is organized as follows. In Section 2 and 3, we give precise definitions for stepped planes, functional stepped surfaces, their codings and review their basic properties. Section 4 is devoted to the generation of a functional stepped surface by a locally finite sequence of flips performed on a given stepped plane. Generalized substitutions associated with a unimodular substitution are introduced in Section 5.1; we prove the image of a stepped plane by such a substitution is again a stepped plane, whose parameters can be explicitly computed. Finally, in Section 5.2, we prove generalized substitutions act on the set of functional stepped surfaces, furthermore the main result of

the present paper, that is, the image of a functional stepped surface is again a functional stepped surface.

We remark that we deal here with three types of objects: functional stepped surfaces, lozenge tilings of the plane and two-dimensional words. There is a straightforward relation between these objects: there is a one-to-one correspondence between lozenge tilings and functional stepped surfaces containing the origin, or functional stepped surfaces up to a translation by a multiple of the diagonal vector  $(1, 1, 1)$  (of course, the translate of a stepped surface by this vector gives the same lozenge tiling by projection): any tiling can be lifted in a unique way, up to translation, to a functional stepped surface, as it is intuitively clear by looking at the picture of a tiling (see for instance Figure 3). The map which associates with a lozenge tiling the corresponding symbolic coding is obviously one-to-one, but not onto; the set of words obtained in this way can be completely described by a local condition (see [Jam04,JP05]). Hence the multidimensional substitutions we deal with here can be equivalently defined as acting either on functional stepped surfaces, or on their codings as a two-dimensional word over a three-letter alphabet, or lastly, on the corresponding tiling of the plane by lozenges. For the sake of clarity, we choose here to focus on the first point of view, that is, on multidimensional substitutions acting on faces of functional stepped surfaces.

## 2 Stepped planes

There are several ways to approximate planes by integer points, see for instance the survey [BCK04]. All these methods boil down to selecting integer points within a bounded distance from the considered plane and such objects are called *discrete planes*. In the present paper, we deal with an approach inspired by the formalism of [AI01], see also [IO93,IO94,BV00,ABI02].

Let  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  denote the canonical basis of  $\mathbb{R}^3$ . Let  $\mathbf{x} \in \mathbb{Z}^3$  and  $i \in \{1, 2, 3\}$ . The *face*  $(\mathbf{x}, i^*)$  is the subset of  $\mathbb{R}^3$  defined as follows:

$$(\mathbf{x}, i^*) = \left\{ \mathbf{x} + \sum_{j \neq i} \lambda_j \mathbf{e}_j, \lambda_j \in [0, 1] \right\}.$$

The integer  $i \in \{1, 2, 3\}$  is called the *type* of the face  $(\mathbf{x}, i^*)$ . We denote by  $\mathfrak{F}$  the set of faces  $\mathfrak{F} = \{(\mathbf{x}, i^*), \mathbf{x} \in \mathbb{Z}^3, i \in \{1, 2, 3\}\}$ , and by  $\mathcal{G}$ , the set of (finite or infinite) unions of faces of  $\mathfrak{F}$ . Endowed with the union operation,  $\mathcal{G}$  is a monoid. In the sequel of the present paper, we need more structure on the set  $\mathcal{G}$ . We provide  $\mathcal{G}$  with a distance as follows:

**Definition 1 (Distance between sets of faces)** *Given  $\mathcal{E}$  and  $\mathcal{E}'$  in  $\mathcal{G}$ , we*




Figure 1. Examples of faces in  $\mathbb{R}^3$ .

set:

$$d(\mathcal{E}, \mathcal{E}') = 2^{-\sup\{r \in \mathbb{R}_+ \cup \{\infty\}, \forall (\mathbf{v}, i^*) \in \mathfrak{F}, \|\mathbf{v}\|_\infty \leq r, \text{ then } (\mathbf{v}, i^*) \subseteq \mathcal{E} \iff (\mathbf{v}, i^*) \subseteq \mathcal{E}'\}}.$$

One easily checks that  $d : \mathcal{G} \times \mathcal{G} \rightarrow [0, 1]$  defines a distance on the set  $\mathcal{G}$ . Roughly speaking, the larger the balls  $B(0, r) = \{\mathbf{x} \in \mathbb{R}^3, \|\mathbf{x}\|_\infty < r\}$  the sets  $\mathcal{E}$  and  $\mathcal{E}'$  coincide on, the closer the sets  $\mathcal{E}$  and  $\mathcal{E}'$  are. In all that follows,  $\mathcal{G}$  stands for the union  $\mathcal{G}$  provided with the topology induced by the distance  $d$ .

From now on, we denote by  $\mathbb{R}_+^3$  the set of vectors in  $\mathbb{R}^3$  with positive coordinates. We then define *stepped planes* as a particular set of faces as follows:

**Definition 2 (Stepped plane)** Let  $\mathbf{v} \in \mathbb{R}_+^3$  and  $\mu \in \mathbb{R}$ . The stepped plane with normal vector  $\mathbf{v}$  and translation parameter  $\mu$  is the subset  $\mathfrak{P}(\mathbf{v}, \mu)$  of  $\mathcal{G}$  defined as follows:

$$\mathfrak{P}(\mathbf{v}, \mu) = \bigcup_{i=1}^3 \bigcup_{\substack{\mathbf{x} \in \mathbb{Z}^3 \\ 0 \leq \langle \mathbf{x}, \mathbf{v} \rangle + \mu < v_i}} (\mathbf{x}, i^*).$$




Figure 2. A piece of a stepped plane in  $\mathbb{R}^3$ .

In other words, one has:

**Proposition 1 ([IO93, IO94])** Let  $\mathbf{v} \in \mathbb{R}_+^3$  and  $\mu \in \mathbb{R}$ . The stepped plane  $\mathfrak{P}(\mathbf{v}, \mu)$  is the boundary of the union of the unit cubes intersecting the open half-space  $\{\mathbf{x} \in \mathbb{R}^3, \langle \mathbf{v}, \mathbf{x} \rangle + \mu < 0\}$ . The set  $\mathfrak{P}(\mathbf{v}, \mu) \cap \mathbb{Z}^3$  is called the set of vertices of  $\mathfrak{P}(\mathbf{v}, \mu)$ .

Let  $\Delta$  be the diagonal plane of equation  $x_1 + x_2 + x_3 = 0$  and let  $\pi$  be the orthogonal projection onto  $\Delta$ . By abuse of notation, we also denote by  $\pi$  the map  $\pi : \mathbb{Z}^3 \rightarrow \mathbb{Z}^2$  defined by  $\pi(\mathbf{x}) = (x_1 - x_3, x_2 - x_3)$ . By construction, a stepped plane is a union of (closed) faces of type 1, 2 or 3. By a relevant choice of one distinguished vertex for each face, depending on the type of the

latter, we obtain a canonical bijection between the vertices of  $\mathfrak{P}(\mathbf{v}, \mu)$  and the two-dimensional lattice  $\mathbb{Z}^2$ . One thus introduces the map  $v : \mathcal{F} \longrightarrow \mathbb{Z}^3$  defined by  $v(\mathbf{x}, i^*) = \mathbf{x} + \mathbf{e}_1 + \cdots + \mathbf{e}_{i-1}$ , for  $i \in \{1, 2, 3\}$ .

**Proposition 2** ([BV00,ABI02]) *Let  $\mathbf{v} \in \mathbb{R}_+^3$ ,  $\mu \in \mathbb{R}$ . One has*

$$\mathfrak{P}(\mathbf{v}, \mu) \cap \mathbb{Z}^3 = v(\{(\mathbf{x}, i^*), (\mathbf{x}, i^*) \in \mathfrak{P}(\mathbf{v}, \mu)\}),$$

and thus

$$\forall (m_1, m_2) \in \mathbb{Z}^2, \exists! (\mathbf{x}, i^*) \in \mathfrak{P}(\mathbf{v}, \mu), \pi \circ v(\mathbf{x}, i^*) = (m_1, m_2).$$

Furthermore, the restriction of the projection map  $\pi$  to  $\mathfrak{P}(\mathbf{v}, \mu)$  is one-to-one and onto  $\Delta$ ; the projections of the faces of the stepped plane  $\mathfrak{P}(\mathbf{v}, \mu)$  tile the diagonal plane  $\Delta$  with three kinds of lozenges (see Figure 3).




Figure 3. From a stepped plane to a tiling of the plane  $\Delta$  by three kind of lozenges.

Let us recall that according to J.-P. Reveillès' terminology [Rev91], given  $\mathbf{v} \in \mathbb{R}^3$  and  $(\mu, \omega) \in \mathbb{R}^2$ , the *lower* (resp. *upper*) *arithmetic hyperplane*  $\mathfrak{P}(\mathbf{v}, \mu, \omega)$  is defined as the set of points  $\mathbf{x} \in \mathbb{Z}^3$  satisfying  $0 \leq \langle \mathbf{x}, \mathbf{v} \rangle + \mu < \omega$  (resp.  $0 < \langle \mathbf{x}, \mathbf{v} \rangle + \mu \leq \omega$ ). Moreover, if  $\omega = \sum |v_i| = \|\mathbf{v}\|_1$ , then  $\mathfrak{P}(\mathbf{v}, \mu, \omega)$  is said to be *standard* whereas it is said to be *naive* if  $\omega = \max |v_i| = \|\mathbf{v}\|_\infty$ . One thus checks that the set  $\{\mathbf{x} \in \mathbb{Z}^3, \exists i \in \{1, 2, 3\}, (\mathbf{x}, i^*) \in \mathfrak{P}(\mathbf{v}, \mu)\}$  is the naive arithmetic discrete plane  $\mathfrak{P}(\mathbf{v}, \mu, \|\mathbf{v}\|_\infty)$ , whereas  $\mathfrak{P}(\mathbf{v}, \mu) \cap \mathbb{Z}^3$  is the standard arithmetic discrete plane  $\mathfrak{P}(\mathbf{v}, \mu, \|\mathbf{v}\|_1)$ , according to Proposition 2.

The bijection between the faces of  $\mathfrak{P}(\mathbf{v}, \mu)$  and the lattice  $\mathbb{Z}^2$  ensures us, that, given a point  $(m_1, m_2) \in \mathbb{Z}^2$ , there exists one and only one face  $(\mathbf{x}, i^*)$  of  $\mathfrak{P}(\mathbf{v}, \mu)$  such that  $\pi \circ v(\mathbf{x}, i^*) = (m_1, m_2)$  (see Proposition 2). We thus provide each stepped plane with a two-dimensional coding as follows:

**Definition 3 (Two-dimensional coding of a stepped plane)** *Let  $\mathfrak{P}(\mathbf{v}, \mu)$  be a stepped plane with  $\mathbf{v} \in \mathbb{R}_+^3$  and  $\mu \in \mathbb{R}$ . The two-dimensional coding of the stepped plane  $\mathfrak{P}(\mathbf{v}, \mu)$  is the two-dimensional word  $u \in \{1, 2, 3\}^{\mathbb{Z}^2}$  defined by: for all  $(m_1, m_2) \in \mathbb{Z}^2$  and all  $i \in \{1, 2, 3\}$ ,*

$$u_{m_1, m_2} = i \iff \exists (\mathbf{x}, i^*) \in \mathfrak{P}(\mathbf{v}, \mu) \text{ such that } (m_1, m_2) = \pi \circ v(\mathbf{x}, i^*).$$




Figure 4. From a lozenge tiling of  $\Delta$  to a 2D-word.

From Definition 3 and Proposition 2, an easy computation gives:

**Proposition 3** *Let  $\mathbf{v} \in \mathbb{R}_+^3$ ,  $\mu \in \mathbb{R}$  and  $u \in \{1, 2, 3\}^{\mathbb{Z}^2}$  be the two-dimensional coding of the stepped plane  $\mathfrak{P}(\mathbf{v}, \mu)$ . Let  $(m_1, m_2) \in \mathbb{Z}^2$  and  $i \in \{1, 2, 3\}$ . Then  $u_{m_1, m_2} = i$  if and only if:*

$$m_1 v_1 + m_2 v_2 + \mu \bmod v_1 + v_2 + v_3 \in [v_1 + \dots + v_{i-1}, v_1 + \dots + v_i[.$$

Of course not all the two-dimensional words over the three-letter alphabet  $\{1, 2, 3\}$  code a stepped plane. For instance, a word containing two consecutive 1's and two consecutive 2's in the same row cannot be the two-dimensional coding of a stepped plane. More generally, not all the two-dimensional words over the three-letter alphabet  $\{1, 2, 3\}$  code a lozenge tiling of the plane  $\Delta$ . In order to generalize the notion of stepped plane to the one of functional stepped surface (see Section 3), we use a slightly more precise property of the restriction of the projection map  $\pi$  to  $\mathfrak{P}(\mathbf{v}, \mu)$ .

**Proposition 4** *The restriction of the map  $\pi$  to  $\mathfrak{P}(\mathbf{v}, \mu)$  is a homeomorphism onto the plane  $\Delta$ .*

PROOF. We already know from Proposition 2 that the restriction of  $\pi$  is a bijection. We first note that  $\mathfrak{P}(\mathbf{v}, \mu)$  is a closed subset of  $\mathbb{R}^3$ . This is a direct consequence of the fact that the faces  $(\mathbf{x}, i^*)$  are closed subsets of  $\mathbb{R}^3$  and that each ball  $B(\mathbf{y}, R) = \{\mathbf{y} \in \mathbb{R}^3, \|\mathbf{y} - \mathbf{x}\|_\infty < R\}$ , with  $R > 0$ , intersects a finite number of faces of  $\mathfrak{P}(\mathbf{v}, \mu)$ . Second, since the map  $\pi : \mathbb{R}^3 \longrightarrow \{\mathbf{x} \in \mathbb{R}^3, x_1 + x_2 + x_3 = 0\}$  is closed, then the image of a closed subset of  $\mathbb{R}^3$  is a closed subset of  $\Delta$ . This holds in particular for the closed subsets of  $\mathfrak{P}(\mathbf{v}, \mu)$  and  $\pi^{-1} : \Delta \longrightarrow \mathfrak{P}(\mathbf{v}, \mu)$  is continuous. It follows that the map  $\pi : \mathfrak{P}(\mathbf{v}, \mu) \longrightarrow \Delta$  is a homeomorphism. ■

### 3 Functional stepped surface

It is natural to try to extend the previous definitions and results to more general objects:

**Definition 4** ([Jam04]) *A union  $\mathfrak{S}$  of faces  $(\mathbf{x}, i^*)$ , where  $\mathbf{x} \in \mathbb{Z}^3$  and  $i \in \{1, 2, 3\}$ , is called a functional stepped surface if the restriction of the projection map  $\pi$  to  $\mathfrak{S}$  is a homeomorphism.*

In particular, a stepped plane is a functional stepped surface, according to Proposition 4.

**Proposition 5** ([Jam04,JP05]) *Let  $\mathfrak{S}$  be a functional stepped surface. One has  $\mathfrak{S} \cap \mathbb{Z}^3 = \pi \circ v(\{(\mathbf{x}, i^*), (\mathbf{x}, i^*) \subset \mathfrak{S}\})$ . Furthermore, given  $(m_1, m_2) \in \mathbb{Z}^2$ , there exists a unique face  $(\mathbf{x}, i^*) \subset \mathfrak{S}$  such that  $(m_1, m_2) = \pi \circ v(\mathbf{x}, i^*)$ .*

PROOF. The proof is deduced from a simple case study. ■

The following coding is thus well-defined:

**Definition 5** *A two-dimensional word  $u \in \{1, 2, 3\}^{\mathbb{Z}^2}$  is said to be the coding of the functional stepped surface  $\mathfrak{S}$  if for all  $(m_1, m_2) \in \mathbb{Z}^2$  and for every  $i \in \{1, 2, 3\}$ :  $u_{m_1, m_2} = i \iff \exists (\mathbf{x}, i^*) \in \mathfrak{S}$  such that  $(m_1, m_2) = \pi \circ v(\mathbf{x}, i^*)$ .*

**Proposition 6** *A union of faces  $\mathfrak{S} \subset \mathcal{G}$  is a functional stepped surface if and only if the restriction of  $\pi$  to  $\mathfrak{S}$  is a bijection onto  $\Delta$ .*

PROOF. Let  $\mathfrak{S}$  be a union of faces such that the restriction of  $\pi$  to  $\mathfrak{S}$  is a bijection onto  $\Delta$ . Then,  $\mathfrak{S}$  is a closed set. We just conclude similarly as in the proof of Proposition 4. ■




Figure 5. A stepped surface.

**Proposition 7** Let  $\mathfrak{S}$  and  $\mathfrak{S}'$  be two stepped surfaces. Then one has:

$$\mathfrak{S} = \mathfrak{S}' \quad \Leftrightarrow \quad \mathfrak{S} \cap \mathbb{Z}^3 = \mathfrak{S}' \cap \mathbb{Z}^3.$$

PROOF. Let  $\mathfrak{S}$  be a functional stepped surface. It is sufficient to prove that given the four vertices of a face, say  $(\mathbf{x}, i^*) \in \mathfrak{S}$ , then the whole face is included in  $\mathfrak{S}$ . On the contrary, since the restriction on  $\Delta$  of  $\pi^{-1}$  is continuous, then there exists  $\lambda \neq 0 \in \mathbb{R}$  such that  $(\mathbf{x} + \lambda(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3), i^*) \subset \mathfrak{S}$ . But then, the vertices of  $(\mathbf{x}, i^*)$  and of  $(\mathbf{x} + \lambda(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3), i^*)$  have the same projection by  $\pi$ , hence a contradiction.  $\blacksquare$

**Definition 6** (i) Let  $\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{Z}^3$  and  $\mathbf{x}' = (x'_1, x'_2, x'_3) \in \mathbb{Z}^3$  such that  $\pi(\mathbf{x}) = \pi(\mathbf{x}')$ . We say that  $\mathbf{x}$  is above  $\mathbf{x}'$  if  $x_1 + x_2 + x_3 \geq x'_1 + x'_2 + x'_3$ , otherwise we say that  $\mathbf{x}$  is below  $\mathbf{x}'$ .

(ii) We then say that a functional stepped surface  $\mathfrak{S}$  is above (resp. below) a stepped surface  $\mathfrak{S}'$  if, for any  $\mathbf{x} \in \mathfrak{S} \cap \mathbb{Z}^3$  and  $\mathbf{x}' \in \mathfrak{S}' \cap \mathbb{Z}^3$  such that  $\pi(\mathbf{x}) = \pi(\mathbf{x}')$ ,  $\mathbf{x}$  is above (resp. below)  $\mathbf{x}'$ .

**Definition 7** For  $\mathbf{s} \in \mathbb{Z}^3$ , we define two specific functional stepped surfaces  $\hat{\mathcal{C}}_{\mathbf{s}}$  and  $\check{\mathcal{C}}_{\mathbf{s}}$  by their intersections with  $\mathbb{Z}^3$ , according to Proposition 7 (see Figure 6):

$$\hat{\mathcal{C}}_{\mathbf{s}} \cap \mathbb{Z}^3 = \{\mathbf{s}' \in \mathbb{Z}^3, (s_1 - s'_1)(s_2 - s'_2)(s_3 - s'_3) \neq 0 \text{ and } s'_i \leq s_i, i \in \{1, 2, 3\}\},$$

$$\check{\mathcal{C}}_{\mathbf{s}} \cap \mathbb{Z}^3 = \{\mathbf{s}' \in \mathbb{Z}^3, (s_1 - s'_1)(s_2 - s'_2)(s_3 - s'_3) \neq 0 \text{ and } s'_i \geq s_i, i \in \{1, 2, 3\}\}.$$




Figure 6. The stepped surfaces  $\hat{\mathcal{C}}_{\mathbf{s}}$  (left) and  $\check{\mathcal{C}}_{\mathbf{s}}$  (right).

**Proposition 8** Let  $\mathfrak{S}$  be a stepped surface and let  $\mathbf{s} \in \mathfrak{S} \cap \mathbb{Z}^3$ . Then  $\hat{\mathcal{C}}_{\mathbf{s}}$  (resp.  $\check{\mathcal{C}}_{\mathbf{s}}$ ) is below (resp. above)  $\mathfrak{S}$ .

PROOF. Let  $\mathfrak{S}$  be a stepped surface,  $\mathbf{s} \in \mathfrak{S} \cap \mathbb{Z}^3$  and  $\mathbf{s}' \in \check{\mathcal{C}}_{\mathbf{s}}$ . Let us assume that  $s'_3 = s_3$ . The other cases can be handled similarly. We introduce a finite sequence of points  $(\mathbf{w}_k)_{0 \leq k \leq s_1 - s'_1 + s_2 - s'_2}$  with values in  $\check{\mathcal{C}}_{\mathbf{s}}$  defined as follows:  $\mathbf{w}_0 = \mathbf{s}$ ; for  $1 \leq k \leq s_1 - s'_2$ ,  $\mathbf{w}_k = \mathbf{s} + k\mathbf{e}_1$ ; and for  $s_1 - s'_1 + 1 \leq k \leq s_1 - s'_1 + s_2 - s'_2$ ,  $\mathbf{w}_k = \mathbf{s} + (s'_1 + s_1)\mathbf{e}_1 + k\mathbf{e}_2$ . One has  $\mathbf{w}_{s_1 - s'_1 + s_2 - s'_2} = \mathbf{s}$ . We

now denote by  $(\mathbf{w}'_k)_{0 \leq k \leq s_1 - s'_1 + s_2 - s'_2}$  the sequence of points with values in  $\mathfrak{S}$  such that  $\pi(\mathbf{w}'_k) = \pi(\mathbf{w}_k)$ . In particular,  $\pi(\mathbf{w}'_{s_1 - s'_1 + s_2 - s'_2}) = \pi(\mathbf{s}')$ . One has, for  $0 \leq k \leq s_1 - s'_1$ ,  $\mathbf{w}_{k+1} - \mathbf{w}_k = \mathbf{e}_1$ . The functional stepped surface  $\mathfrak{S}$  is a connected subset of  $\mathbb{R}^3$ ; indeed, it is the image of a connected set, namely  $\Delta$ , by a continuous map. Hence,  $\mathbf{w}_{k+1} - \mathbf{w}'_{k+1} \in \{\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3, 0\}$ . Similarly, for  $s_1 - s'_1 + 1 \leq k \leq s_1 - s'_1 + s_2 - s'_2$ , one has  $\mathbf{w}_{k+1} - \mathbf{w}_k = \mathbf{e}_2$ , which yields  $\mathbf{w}_{k+1} - \mathbf{w}'_{k+1} \in \{\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3, 0\}$ . One thus gets that  $\mathbf{s}'$  is above  $\mathbf{w}'_{s_1 - s'_1 + s_2 - s'_2}$ . We similarly prove that  $\hat{\mathcal{C}}_{\mathbf{s}}$  is below  $\mathfrak{S}$ .  $\blacksquare$

**Definition 8** *A lozenge tiling of  $\Delta$  is defined as a union of lozenges  $\pi(\mathbf{x}, i^*)$ , for  $(\mathbf{x}, i^*) \in \mathcal{F}$  such that this union covers  $\Delta$ , and furthermore, the interiors of two lozenges do not intersect.*

**Theorem 9** *Let  $\mathcal{D} = \bigcup_{(\mathbf{x}, i^*) \in \mathcal{T}} \pi(\mathbf{x}, i^*)$ , where  $\mathcal{T} \subset \mathcal{F}$ , be a lozenge tiling of  $\Delta$ . Then there exists a unique functional stepped surface  $\mathfrak{S}$ , up to translation by the vector  $\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$ , of the form  $\bigcup_{(\mathbf{x}, i^*) \in \mathcal{T}} \pi(\mathbf{y}(\mathbf{x}), i^*)$  with  $(\mathbf{y}(\mathbf{x}), i^*) \in \mathcal{F}$ , and  $\pi(\mathbf{x}, i^*) = \pi(\mathbf{y}(\mathbf{x}), i^*)$ , for all  $(\mathbf{x}, i^*) \in \mathcal{T}$ . Such a functional stepped surface is said to project onto  $\mathcal{D}$ .*

**PROOF.** Let  $\mathcal{D} = \bigcup_{(\mathbf{x}, i^*) \in \mathcal{T}} \pi(\mathbf{x}, i^*)$ , where  $\mathcal{T} \subset \mathcal{F}$ , be a lozenge tiling of  $\Delta$ . Let us note that there is no reason for the union of faces  $\bigcup_{(\mathbf{x}, i^*) \in \mathcal{T}} \pi(\mathbf{x}, i^*)$  to be a functional stepped surface.

Let  $\Gamma$  be the lattice of  $\Delta$  generated by the vectors  $\pi(\mathbf{e}_1)$ ,  $\pi(\mathbf{e}_2)$ , and  $\pi(\mathbf{e}_3)$ . Similarly as in the proof of Proposition 5 (see e.g. [Jam04,JP05]), one proves by a finite case study that the points of  $\mathcal{D} \cap \Gamma = \Gamma$  are exactly the vertices of the lozenges  $\pi(\mathbf{x}, i^*)$ , for  $(\mathbf{x}, i^*) \in \mathcal{T}$ , and are also exactly the points  $\pi \circ v(\mathbf{x}, i^*)$ , for  $(\mathbf{x}, i^*) \in \mathcal{T}$ . In other words, we have chosen a distinguished vertex for each lozenge  $\pi(\mathbf{x}, i^*)$ : for any  $\gamma \in \Gamma$ , there exists a unique  $i_\gamma \in \{1, 2, 3\}$  such that  $i_\gamma^*$  is the type of the lozenge whose distinguished vertex is  $\gamma$ . One thus gets  $\mathcal{D} = \bigcup_{\gamma \in \Gamma} \pi(\gamma, i_\gamma^*)$ . Furthermore, there is a one-to-one correspondence between the faces  $(\mathbf{x}, i^*)$  of  $\mathcal{T}$ , and the faces  $(\gamma, i_\gamma^*)$ , for  $\gamma \in \Gamma$ . Hence a functional stepped surface projects onto  $\mathcal{D}$  if and only if it is of the form  $\bigcup_{\gamma \in \Gamma} \pi(\mathbf{x}_\gamma, i_\gamma^*)$  such that for every  $\gamma \in \Gamma$ , then  $\pi(\mathbf{x}_\gamma) = \gamma$ .

Let us first exhibit a functional stepped surface of the form  $\bigcup_{\gamma \in \Gamma} \pi(\mathbf{x}_\gamma, i_\gamma^*)$  such that for every  $\gamma \in \Gamma$ ,  $\pi(\mathbf{x}_\gamma) = \gamma$ . For that purpose, we introduce the oriented graph  $G = (V, E)$  whose set of vertices is  $V = \Gamma$ , and whose set of edges  $E$  is equal to the set of edges of the lozenges  $\pi(\mathbf{x}, i^*)$ , for  $(\mathbf{x}, i^*) \in \mathcal{T}$ , endowed with both orientations. We first define a *weight* function on the edges of  $G$  as follows: for any  $\gamma, \gamma' \in \Gamma$  such that the oriented edge  $e(\gamma, \gamma')$  from  $\gamma$  to  $\gamma'$  belongs to  $E$ , then one sets  $w(\gamma, \gamma') = 1$ , if  $\gamma' = \gamma + \pi(\mathbf{e}_3)$ ,  $w(\gamma, \gamma') = -1$ , if  $\gamma' = \gamma - \pi(\mathbf{e}_3)$ , and 0, otherwise. One checks by induction on the lengths of the cycles of  $G$  that the sum of the weights of a cycle is equal to zero. We thus can define a *height* function on the vertices of  $G$  as follows: one sets  $h_0 = 0$ ,

and for any  $\gamma, \gamma' \in \Gamma$  such that the edge with vertices  $\gamma$  and  $\gamma'$  belongs to  $E$ , then  $h_{\gamma'} = h_\gamma + 1$ , if  $\gamma' = \gamma + \pi(\mathbf{e}_3)$ ,  $h_{\gamma'} = h_\gamma - 1$ , if  $\gamma' = \gamma - \pi(\mathbf{e}_3)$ , and  $h_{\gamma'} = h_\gamma$ , otherwise. One checks that this function is well-defined for any vertex of  $G$  since the graph  $G$  is connected, and according to the properties of the weight function. We then define for  $\gamma \in \Gamma$ ,  $\mathbf{x}_\gamma$  as the point of  $\mathbb{R}^3$  equal to  $\gamma + h_\gamma(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3)$ , and  $i_\gamma^*$  as the type of the unique lozenge whose distinguished vertex has coordinates  $\gamma$ . We now consider  $\mathfrak{S} = \bigcup_{\gamma \in \Gamma} (\mathbf{x}_\gamma, i_\gamma^*)$ . It remains to prove that  $\mathfrak{S}$  is a functional stepped surface. According to Proposition 6, this is a direct consequence of the fact that the restriction of  $\pi$  to  $\mathfrak{S}$  is a bijection, which can be deduced from its construction.

Let us consider now a functional stepped surface that contains the origin 0 of  $\mathbb{R}^3$  and that projects onto  $\mathcal{D}$ ; it is of the form  $\bigcup_{\gamma \in \Gamma} (\mathbf{y}_\gamma, i_\gamma^*)$  with  $\mathbf{y}_\gamma - \gamma \in \mathbb{Z}(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3)$ , for all  $\gamma \in \Gamma$ . A functional stepped surface is connected, hence one checks that necessarily,  $\mathbf{y}_\gamma = \gamma + h_\gamma$ . ■

#### 4 Flips acting on stepped surfaces

Let us define, for  $\mathbf{s} \in \mathbb{Z}^3$ , two specific unions of faces (see Figure 7):

$$\check{c}_\mathbf{s} = \bigcup_{i=1}^3 (\mathbf{s}, i^*) \quad \text{and} \quad \hat{c}_\mathbf{s} = \bigcup_{i=1}^3 (\mathbf{s} + \mathbf{e}_i, i^*).$$

Let us note that a functional stepped surface cannot contain simultaneously  $\hat{c}_\mathbf{s}$  and  $\check{c}_\mathbf{s}$ ; furthermore, if a functional stepped surface contains one of them, then by exchanging both unions, we still have a functional stepped surface. This leads us to define a simple operation on functional stepped surfaces, the so-called *flip*, such as depicted in Figure 7:

**Definition 9 (Flip)** *Let  $\mathbf{s} \in \mathbb{Z}^3$ . The flip map  $\varphi_\mathbf{s} : \mathcal{G} \rightarrow \mathcal{G}$  is defined as follows: if a union of faces  $\mathcal{E} \in \mathcal{G}$  contains  $\hat{c}_\mathbf{s}$  (resp.  $\check{c}_\mathbf{s}$ ), then  $\varphi_\mathbf{s}(\mathcal{E})$  is obtained by replacing  $\hat{c}_\mathbf{s}$  by  $\check{c}_\mathbf{s}$  (resp.  $\check{c}_\mathbf{s}$  by  $\hat{c}_\mathbf{s}$ ); otherwise,  $\varphi_\mathbf{s}(\mathcal{E}) = \mathcal{E}$ .*




Figure 7. The action of flip  $\varphi_\mathbf{s}$ , for  $\mathbf{x} \in \mathbb{Z}^3$ :  $\check{c}_\mathbf{s}$  (left) is exchanged with  $\hat{c}_\mathbf{s}$  (right).

According to Theorem 9, we can perform a flip on a functional stepped surface if and only if one can perform a classic flip in the sense, e.g., of [Thu89] on the

lozenge tiling of the plane which corresponds to this functional stepped surface.

We are now interested in performing on a functional stepped surface, not only one flip, but a sequence of flips. We first need to introduce the following notion:

**Definition 10 (Locally finiteness)** *A sequence of flips  $(\varphi_{\mathbf{s}_n})_{n \in \mathbb{N}}$  is said to be locally finite if, for any  $n_0 \in \mathbb{N}$ , the set  $\{\mathbf{s}_n \in \mathbb{Z}^3, \pi(\mathbf{s}_n) = \pi(\mathbf{s}_{n_0})\}$  is bounded.*

Then one has:

**Proposition 10** *Let  $\mathfrak{S}$  be a functional stepped surface and  $(\varphi_{\mathbf{s}_n})_{n \in \mathbb{N}^*}$  be a locally finite sequence of flips such that the following limit exists:*

$$\mathfrak{S}' = \lim_{n \rightarrow \infty} \varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S}).$$

*Then,  $\mathfrak{S}'$  is a stepped surface.*

**PROOF.** By performing a single flip on a stepped surface, one easily checks that one obtains a union of faces still homeomorphic by  $\pi$  to  $\Delta$ , i.e., a functional stepped surface. The case of the action of a finite number of flips is straightforward. Suppose now we perform a locally finite sequence of flips  $(\varphi_{\mathbf{s}_n})_{n \in \mathbb{N}}$  on the functional stepped surface  $\mathfrak{S}$  such that  $(\varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S}))_{n \in \mathbb{N}}$  is convergent in  $\mathcal{G}$ . According to Proposition 6, it is sufficient to prove that the restriction of  $\pi$  to  $\mathfrak{S}'$  is a bijection onto  $\Delta$ . Let  $\mathbf{x}$  and  $\mathbf{y}$  be two points of  $\mathfrak{S}'$  such that  $\pi(\mathbf{x}) = \pi(\mathbf{y})$ . There exists  $n \in \mathbb{N}$  such that  $\mathbf{x}, \mathbf{y} \in \varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S})$ . Since  $\varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S})$  is a functional stepped surface, it follows that  $\mathbf{x} = \mathbf{y}$ . We thus have proved that the restriction of  $\pi$  is one-to-one.

Let  $\mathbf{z} \in \Delta$ . Let  $A$  be a bounded subset of  $\Delta$  containing  $\mathbf{x}$ . By local finiteness of the sequence  $(\varphi_{\mathbf{s}_n})_{n \in \mathbb{N}^*}$ , there exists  $n_0 \in \mathbb{N}$  such that, if  $n \geq n_0$ , then  $\pi(\mathbf{s}_n) \notin A$ . Take  $n_1 \geq n_0$ ; we also assume  $n_1$  large enough such that  $\mathfrak{S}'$  and  $\varphi_{\mathbf{s}_{n_1}} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S})$  coincide on their intersection with  $\pi^{-1}(A)$ . Let  $\mathbf{y} \in \varphi_{\mathbf{s}_{n_1}} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S})$  such that  $\pi(\mathbf{y}) = \mathbf{z}$ . Then one has  $\mathbf{y} \in \varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S})$  for all  $n \geq n_1$ , and thus  $\mathbf{y} \in \mathfrak{S}'$ . We have proved that the restriction of  $\pi$  is onto, which concludes the proof.  $\blacksquare$

Thus, flips allow to transform functional stepped surfaces into functional stepped surfaces. However, one cannot necessarily transform a *given* functional stepped surface into another *given* one by a locally finite sequence of flips. See Figure 8 for some examples of (un)accessibility by flips.

In order to characterize the (un)accessibility by flips between stepped surfaces, we introduce the notion of *shadows*, illustrated in Figure 9:

**Definition 11 (Shadows)** *Let  $\mathfrak{S}$  be a functional stepped surface. We define*




Figure 8. One can transform the first stepped surface into the second one, and conversely, by performing a finite number of flips. A locally finite sequence of flips allows one to transform the second stepped surface into the third one (we perform an infinite and locally finite sequence of flips which rejects to the infinity the only face of type 1\*), but the converse transformation is impossible (no flip can be performed). Lastly, we can neither transform by flips the fourth stepped surface into the third one, nor conversely.

three projection maps from  $\mathbb{R}^3$  to  $\mathbb{R}^2$  by:

$$\pi_1 : (x_1, x_2, x_3) \mapsto (x_2, x_3) \quad \pi_2 : (x_1, x_2, x_3) \mapsto (x_1, x_3)$$

and

$$\pi_3 : (x_1, x_2, x_3) \mapsto (x_1, x_2).$$

The shadows of  $\mathfrak{S}$  are respectively defined as the three images of the stepped surface  $\mathfrak{S}$  by these maps.




Figure 9. The shadows of the stepped surfaces of Figure 8. The central stepped surface has all its shadows included in the corresponding ones of the leftmost stepped surface. The shadows of the rightmost stepped surface are neither included in the shadow of the other stepped surfaces, nor contain them.

Considering the functional stepped surfaces of Figure 8, it is worth remarking that one can transform one functional stepped surface into another one if and only if the shadows of the first one contain the respective shadows of the second one (see Figure 9). This turns out to be a general fact:

**Proposition 11** *Let  $\mathfrak{S}$  and  $\mathfrak{S}'$  be two functional stepped surfaces. The following assertions are equivalent:*

(i) *There exists a locally finite sequence  $(\varphi_{\mathbf{s}_n})_{n \in \mathbb{N}}$  of flips such that*

$$\mathfrak{S}' = \lim_{n \rightarrow \infty} \varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{S});$$

(ii) *the three shadows of  $\mathfrak{S}'$  are included in the corresponding shadows of  $\mathfrak{S}$ .*

**PROOF.** Since  $\hat{c}_{\mathfrak{S}}$  and  $\check{c}_{\mathfrak{S}}$  have the same shadows, performing a flip does not modify the shadows of a functional stepped surface. By performing a *sequence*

of flips, the shadows cannot be extended. However, note that they can be reduced (recall the example of Figure 9). Thus, a stepped surface  $\mathfrak{S}'$  can be obtained by performing a locally finite sequence of flips on a stepped surface  $\mathfrak{S}$  only if the three shadows of  $\mathfrak{S}'$  are included in the corresponding shadows of  $\mathfrak{S}$ .

Conversely, let  $\mathfrak{S}'$  and  $\mathfrak{S}$  be two functional stepped surfaces such that the three shadows of  $\mathfrak{S}'$  are included in the corresponding shadows of  $\mathfrak{S}$ . Let us consider a vertex  $\mathbf{x} \in \mathfrak{S}' \cap \mathbb{Z}^3$  of the functional stepped surface  $\mathfrak{S}'$ . With no loss of generality, we can suppose that  $\mathbf{x}$  is above the stepped surface  $\mathfrak{S}$ , according to Definition 6. We associate with this vertex  $\mathbf{x} \in \mathfrak{S}'$  the following union of faces of  $\mathfrak{S}$  (see Figure 10):

$$\mathcal{T}_{\mathbf{x}} = \bigcup_{\substack{(\mathbf{x}', i^*) \subset \mathfrak{S} \\ x'_i < x_i}} (\mathbf{x}', i^*).$$

Let us prove that  $\mathcal{T}_{\mathbf{x}}$  is a *finite* union of faces. By assumption, the shadow




Figure 10. Given a stepped surface  $\mathfrak{S}$ , a vertex  $\mathbf{x} \in \mathfrak{S}' \cap \mathbb{Z}^3$  defines a subset  $\mathcal{T}_{\mathbf{x}}$  of  $\mathfrak{S}$ . Left: the vertex  $\mathbf{x}$  and  $\mathbf{x}_3$ , and the boundaries of  $\mathcal{T}_{\mathbf{x}}$ . Right: the faces of  $\mathcal{T}_{\mathbf{x}}$  (in white). To  $\mathcal{T}_{\mathbf{x}}$  corresponds a lozenge tiling of a bounded and simply connected domain of  $\mathbb{R}^2$ .

$\pi_1(\mathfrak{S}')$  is included in the shadow  $\pi_1(\mathfrak{S})$ . In particular,  $\pi_1(\mathbf{x}) \in \pi_1(\mathfrak{S})$ : there exists  $x'_1 \in \mathbb{Z}$  such that  $(x'_1, x_2, x_3) \in \mathfrak{S}$ . Then, according to Proposition 8,  $\mathcal{T}_{\mathbf{x}} \subset \mathfrak{S}$  is above  $\hat{\mathcal{C}}_{(x'_1, x_2, x_3)}$ . Since  $\mathbf{x}'' \in \mathcal{T}_{\mathbf{x}}$  satisfies  $x''_2 < x_2$  and  $x''_3 < x_3$ , this yields that  $x''_1 \geq x'_1$ . Similarly, there exist  $x'_2 \in \mathbb{Z}$  and  $x'_3 \in \mathbb{Z}$  such that if  $\mathbf{x}'' \in \mathcal{T}_{\mathbf{x}}$ , then  $x_2 > x''_2$  and  $x_3 > x''_3$ . Finally,  $\mathbf{x}'' \in \mathcal{T}_{\mathbf{x}}$  yields  $x'_1 \leq x''_1 \leq x_1$ ,  $x'_2 \leq x''_2 \leq x_2$  and  $x'_3 \leq x''_3 \leq x_3$ . Thus,  $\mathcal{T}_{\mathbf{x}}$  is bounded, i.e., it is a finite union of faces.

Let us now consider the union of faces  $\hat{\mathcal{T}}_{\mathbf{x}}$  which is included in  $\hat{\mathcal{C}}_{\mathbf{x}}$  and satisfies  $\pi(\hat{\mathcal{T}}_{\mathbf{x}}) = \pi(\mathcal{T}_{\mathbf{x}})$  (see Figure 11, left). Similarly as  $\mathcal{T}_{\mathbf{x}}$ ,  $\hat{\mathcal{T}}_{\mathbf{x}}$  is a finite union of faces. A classic result of the theory of lozenge tilings (see, e.g., [Thu89]) yields that the tiling corresponding to  $\mathcal{T}_{\mathbf{x}}$  can be transformed by performing a finite number of flips into the tiling corresponding to  $\hat{\mathcal{T}}_{\mathbf{x}}$ . In terms of stepped surfaces, this means that a finite number of flips transforms  $\mathcal{S}$  (which contains  $\mathcal{T}_{\mathbf{x}}$ ) into a stepped surface which contains  $\hat{\mathcal{T}}_{\mathbf{x}}$ , hence the vertex  $\mathbf{x}$  of  $\mathcal{S}'$  (since  $\mathbf{x} \in \hat{\mathcal{T}}_{\mathbf{x}}$ ) too.

Now, we would like to perform such a finite number of flips for each  $\mathbf{x} \in \mathfrak{S}' \cap \mathbb{Z}^3$




Figure 11. By performing a finite number of flips, one transforms  $\mathcal{T}_{\mathbf{x}}$  (Figure 10, right) into the union of faces  $\hat{\mathcal{T}}_{\mathbf{x}}$  (left, with white faces). We obtain a stepped surface which contains the vertex  $\mathbf{x}$  of  $\mathfrak{S}'$ , similarly as  $\hat{\mathcal{T}}_{\mathbf{x}}$  does (right). By performing such a finite number of flips for each vertex of  $\mathfrak{S}'$ , this transforms the stepped surface  $\mathfrak{S}$  into the stepped surface  $\mathfrak{S}'$ .

$\mathbb{Z}^3$ , in order to transform by an infinite sequence of flips the functional stepped surface  $\mathfrak{S}$  into a functional stepped surface which would contain all the vertices  $\mathfrak{S}' \cap \mathbb{Z}^3$ , i.e., into  $\mathfrak{S}'$ , by Proposition 7. The only one problem could be the following one: by performing the flips to obtain a stepped surface containing a given  $\mathbf{x}$  in  $\mathfrak{S}' \cap \mathbb{Z}^3$ , we could lose a vertex  $\mathbf{x}'$  of  $\mathfrak{S}' \cap \mathbb{Z}^3$  previously obtained by performing flips. However, the flips performed to obtain  $\mathbf{x} \in \mathfrak{S}' \cap \mathbb{Z}^3$  are performed *below*  $\hat{\mathcal{T}}_{\mathbf{x}}$ , in particular *below*  $\mathfrak{S}'$  since  $\hat{\mathcal{T}}_{\mathbf{x}} \subset \hat{\mathcal{C}}_{\mathbf{x}}$  and  $\hat{\mathcal{C}}_{\mathbf{x}}$  is below  $\mathfrak{S}'$  by Proposition 8. Hence, we do not lose the previously obtained vertices of  $\mathfrak{S}' \cap \mathbb{Z}^3$ , and the whole (infinite) sequence of flips thus transforms  $\mathfrak{S}$  into  $\mathfrak{S}'$ .

To conclude, we note that the finite number of flips performed to obtain a stepped surface containing a vertex  $\mathbf{x}$  of  $\mathfrak{S}' \cap \mathbb{Z}^3$  are performed at a *bounded* distance from  $\mathbf{x}$ . This yields that the previous sequence of flips (that is, the one used to obtain the stepped surface containing *all* the vertices of  $\mathfrak{S}' \cap \mathbb{Z}^3$ ) contains, for each  $\pi(\mathbf{x}) \in \pi(\mathfrak{S}' \cap \mathbb{Z}^3) = \pi(\mathbb{Z}^3)$ , a *finite* number of flips  $\varphi_{\mathbf{x}'}$  such that  $\pi(\mathbf{x}') = \pi(\mathbf{x})$ . Thus, this is a *locally finite* sequence of flips. This completes the proof. ■

Hence, flips transform functional stepped surfaces into functional stepped surfaces, and we have obtained a necessary and sufficient condition - in terms of shadows - under which a given functional stepped surface can be transformed by flips into another one. In particular, we can use these results to give an equivalent definition of functional stepped surfaces:

**Theorem 12** *A union of faces  $\mathfrak{U} \in \mathcal{G}$  is a functional stepped surface if and only if there exist a stepped plane  $\mathfrak{P}$  and a locally finite sequence of flips  $(\varphi_{\mathbf{s}_n})_{n \in \mathbb{N}}$  such that*

$$\mathfrak{U} = \lim_{n \rightarrow \infty} \varphi_{\mathbf{s}_n} \circ \dots \circ \varphi_{\mathbf{s}_1}(\mathfrak{P}).$$

PROOF. Since a stepped plane is a functional stepped surface, Proposition 10 yields that the limit of a sequence of functional stepped surfaces obtained by performing a locally finite sequence of flips over a stepped plane is a functional stepped surface. Conversely, it is easy to check that the three shadows of a stepped plane with normal vector  $\mathbf{v} \in \mathbb{R}_+^3$  such that  $v_1 v_2 v_3 \neq 0$  are equal to the whole plane  $\mathbb{R}^2$ . Therefore, according to Proposition 11, one can transform by flips any stepped plane  $\mathfrak{P}$  into a given stepped surface  $\mathfrak{S}$ . ■

## 5 Generalized substitutions

We first review in Section 5.1 the notion of *generalized substitutions* [AI01]; we then discuss in Section 5.2 the way they act on stepped planes and more generally functional stepped surfaces.

### 5.1 First definitions

Let  $\mathcal{A}$  be a finite alphabet and let  $\mathcal{A}^*$  be the set of finite words over  $\mathcal{A}$ . The empty word is denoted by  $\varepsilon$ . A *substitution* is an endomorphism of the free monoid  $\mathcal{A}^*$  such that the image of every letter of  $\mathcal{A}$  is non-empty. Such a definition naturally extends to infinite or biinfinite words in  $\mathcal{A}^{\mathbb{N}}$  and  $\mathcal{A}^{\mathbb{Z}}$ .

Assume  $\mathcal{A} = \{1, 2, 3\}$  and let  $\sigma$  be a substitution over  $\mathcal{A}$ . The *incidence matrix*  $\mathbf{M}_\sigma$  of  $\sigma$  is the  $3 \times 3$  matrix defined by:

$$\mathbf{M}_\sigma = (|\sigma(j)|_i)_{(i,j) \in \{1,2,3\}^2},$$

where  $|\sigma(j)|_i$  is the number of occurrences of the letter  $i$  in  $\sigma(j)$ .

A substitution  $\sigma$  is then said to be *unimodular* if  $\det \mathbf{M}_\sigma = \pm 1$ . In particular,  $\mathbf{M}_\sigma^{-1}$  has *integer* coefficients. Let  $\mathbf{f} : \{1, 2, 3\}^* \rightarrow \mathbb{N}^3$  be the map defined by  $\mathbf{f}(w) = {}^t(|w|_1, |w|_2, |w|_3)$ . The map  $\mathbf{f}$  is usually called the *Parikh mapping* and is the homomorphism obtained by abelianization of the free monoid  $\mathcal{A}^*$ . One has for every  $w \in \{1, 2, 3\}^*$ ,  $\mathbf{f}(\sigma(w)) = \mathbf{M}_\sigma \cdot \mathbf{f}(w)$ .

**Definition 12 (Generalized substitution [AI01])** *Let  $\sigma$  be a unimodular substitution over  $\{1, 2, 3\}$ . The generalized substitution  $\Theta_\sigma^* : \mathcal{G} \rightarrow \mathcal{G}$  is defined by:*

$$\forall (\mathbf{v}, i^*) \in \mathfrak{F}, \quad \Theta_\sigma^*((\mathbf{v}, i^*)) = \bigcup_{\substack{j, p, s \\ \sigma(j) = p \cdot i \cdot s}} (\mathbf{M}_\sigma^{-1}(\mathbf{v} + \mathbf{f}(s)), j^*).$$

and

$$\forall \mathcal{E} \in \mathcal{G}, \quad \Theta_\sigma^*(\mathcal{E}) = \bigcup_{(\mathbf{v}, i^*) \subseteq \mathcal{E}} \Theta_\sigma^*((\mathbf{v}, i^*)).$$

**Remark 1**

**Example 1** Let  $\sigma : \{1, 2, 3\}^* \longrightarrow \{1, 2, 3\}^*$  be the substitution defined by  $\sigma : 1 \mapsto 13, 2 \mapsto 1, 3 \mapsto 2$ . Then,

$$\mathbf{M}_\sigma = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \mathbf{M}_\sigma^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

This yields (see Figure 12):

$$\begin{aligned} (\mathbf{v}, 1^*) &\mapsto (\mathbf{M}_\sigma^{-1}\mathbf{v} + \mathbf{e}_1 - \mathbf{e}_2, 1^*) \cup (\mathbf{M}_\sigma^{-1}\mathbf{v}, 2^*) \\ \Theta_\sigma^* : (\mathbf{v}, 2^*) &\mapsto (\mathbf{M}_\sigma^{-1}\mathbf{v}, 3^*) \\ (\mathbf{v}, 3^*) &\mapsto (\mathbf{M}_\sigma^{-1}\mathbf{v}, 1^*). \end{aligned}$$




Figure 12. Action of  $\Theta_\sigma^*$  on single faces and on a given union of faces.

There is a natural measure  $\mu$  defined on the elements of  $\mathcal{G}$ , obtained by extension of the two-dimensional Lebesgue measure. Two elements  $\mathcal{E}$  and  $\mathcal{E}'$  of  $\mathcal{G}$  are then said to be  $\mu$ -disjoint if  $\mu(\mathcal{E} \cap \mathcal{E}') = 0$ . In other words, this means that both sets do not intersect, except possibly on edges. A generalized substitution does not necessarily map  $\mu$ -disjoint faces to  $\mu$ -disjoint unions of faces. Consider in Example 1,  $\Theta_\sigma^*(\mathbf{0}, 1^*) \cap \Theta_\sigma^*(\mathbf{e}_3, 3^*)$ . One has  $\Theta_\sigma^*(\mathbf{0}, 1^*) = (\mathbf{e}_1 - \mathbf{e}_2, 1^*) \cup (\mathbf{0}, 2^*)$  and  $\Theta_\sigma^*(\mathbf{e}_3, 3^*) = (\mathbf{e}_1 - \mathbf{e}_2, 1^*)$ , whence  $\mu(\Theta_\sigma^*(\mathbf{0}, 1^*) \cap \Theta_\sigma^*(\mathbf{e}_3, 3^*)) \neq 0$ .

**Definition 13** A generalized substitution  $\Theta_\sigma^*$  is said to act properly on a union of faces  $\mathcal{E} \subset \mathcal{G}$  if  $\mu$ -disjoint faces of  $\mathcal{E}$  are mapped onto  $\mu$ -disjoint unions of faces.

Stepped planes are particularly interesting with respect to this property as shown by Theorem 13 below. Let us assume that the substitution  $\sigma$  is primitive, i.e.,  $\mathbf{M}_\sigma$  admits a power with positive entries. Let  $\mathbf{v} \in \mathbb{R}_+^3$  be a Perron-Frobenius left eigenvector of  $\mathbf{M}_\sigma$  having only positive entries. Then, the generalized substitution  $\Theta_\sigma^*$  is proved in [AI01] to act properly on the stepped plane  $\mathfrak{P}(\mathbf{v}, 0)$ , and to map it onto itself. More generally, one has the following:

**Theorem 13** ([Fer05b]) *Let  $\sigma$  be a unimodular substitution over  $\{1, 2, 3\}$ ,  $\mathbf{v} \in \mathbb{R}_+^3$  and  $\mu \in \mathbb{R}$ . The generalized substitution  $\Theta_\sigma^*$  acts properly on the stepped plane  $\mathfrak{P}(\mathbf{v}, \mu)$ ; furthermore  $\Theta_\sigma^*$  maps  $\mathfrak{P}(\mathbf{v}, \mu)$  onto the stepped plane  $\mathfrak{P}({}^t\mathbf{M}_\sigma\mathbf{v}, \mu)$ .*

## 5.2 Generalized substitutions and functional stepped surfaces

The aim of this section is to extend the previous results to functional stepped surfaces, by proving the main theorem of this paper:

**Theorem 14** *Let  $\sigma$  be a unimodular substitution over  $\{1, 2, 3\}$ . The generalized substitution  $\Theta_\sigma^*$  acts properly on every functional stepped surface. Furthermore, the image by  $\Theta_\sigma^*$  of a functional stepped surface is a functional stepped surface.*

Let us note that a partial version of Theorem 14 has been stated in [ABJ05]. An illustration of Theorem 14 is depicted in Figure 13.




Figure 13. A generalized substitution maps a stepped surface onto a stepped surface.

Several lemmas are required to prove Theorem 14. Let us first prove the continuity of any generalized substitution as a map from  $\mathcal{G}$  to  $\mathcal{G}$  provided with the distance  $d$  (see Definition 1):

**Lemma 15** *Let  $(\mathcal{E}_n)_{n \in \mathbb{N}} \in \mathcal{G}^{\mathbb{N}}$  be a convergent sequence in  $\mathcal{G}$ . Then the se-*

quence  $(\Theta_\sigma^*(\mathcal{E}_n))_{n \in \mathbb{N}} \in \mathcal{G}^{\mathbb{N}}$  is a convergent sequence in  $\mathcal{G}$ . One thus gets:

$$\lim_{n \rightarrow \infty} \Theta_\sigma^*(\mathcal{E}_n) = \Theta_\sigma^*(\lim_{n \rightarrow \infty} \mathcal{E}_n).$$

**PROOF.** Let  $\mathcal{E}$  stand for the limit of the sequence  $(\mathcal{E}_n)_{n \in \mathbb{N}} \in \mathcal{G}^{\mathbb{N}}$ . Let us prove that the sequence  $\Theta_\sigma^*(\mathcal{E}_n)_{n \in \mathbb{N}} \in \mathcal{G}^{\mathbb{N}}$  converges towards  $\Theta_\sigma^*(\mathcal{E})$ . For  $n \in \mathbb{N}$ , let  $r_n$  be such that  $\mathcal{E}_n$  and  $\mathcal{E}$  contain the same faces in a ball of radius  $r \leq r_n$  centered on 0. Let  $M = \max\{\|\mathbf{M}_\sigma \mathbf{f}(s)\|_\infty, s \text{ suffix of } \sigma(i), i \in \{1, 2, 3\}\}$ . Let  $\alpha$  be the modulus of the smallest eigenvalue of  $\mathbf{M}_\sigma^{-1}$ . Let us recall that, for all  $\mathbf{x} \in \mathbb{R}^3$ ,  $\|\mathbf{M}_\sigma^{-1} \mathbf{x}\|_\infty \geq \alpha \|\mathbf{x}\|_\infty$ . Let  $n \in \mathbb{N}$  such that  $r_n > M$  (recall that  $r_n \xrightarrow{n \rightarrow \infty} \infty$ ), and let  $(\mathbf{y}, j^*) \subseteq \Theta_\sigma^*(\mathcal{E}_n)$  such that  $\|\mathbf{y}\|_\infty \leq \alpha(r_n - M)$ . Let  $(\mathbf{x}, i^*) \subseteq \mathcal{E}_n$  such that  $(\mathbf{y}, j^*) \subseteq \Theta_\sigma^*(\mathbf{x}, i^*)$ ; one has  $\mathbf{y} = \mathbf{M}_\sigma^{-1} \mathbf{x} + \mathbf{f}(s)$ , with  $\sigma(j) = p \cdot i \cdot s$ . One deduces  $\|\mathbf{x} + \mathbf{M}_\sigma \mathbf{f}(s)\|_\infty \leq r_n - M$ . Hence  $\|\mathbf{x}\|_\infty \leq r_n$  and  $(\mathbf{y}, j^*) \subseteq \Theta_\sigma^*(\mathcal{E})$ . We show in a similar way that any face  $(\mathbf{y}, j^*)$  included in  $\Theta_\sigma^*(\mathcal{E})$  and satisfying  $\|\mathbf{y}\|_\infty \leq \alpha(r_n - M)$  is included in  $\Theta_\sigma^*(\mathcal{E}_n)$ . In other words,  $d(\Theta_\sigma^*(\mathcal{E}_n), \Theta_\sigma^*(\mathcal{E})) \leq 2^{-\alpha(r_n - M)} \xrightarrow{n \rightarrow \infty} 0$ .  $\blacksquare$

The following lemma plays a key role by relating the action of generalized substitutions to the action of flips, such as depicted in Figure 14:

**Lemma 16** *Let  $\Theta_\sigma^*$  be a generalized substitution that acts properly on  $\mathcal{E} \subset \mathcal{G}$ . Then, for any  $\mathbf{x} \in \mathbb{Z}^3$ ,  $\Theta_\sigma^*$  acts properly on  $\varphi_s(\mathcal{E})$ , and furthermore,  $\Theta_\sigma^*$  maps  $\varphi_s(\mathcal{E})$  onto  $\varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}}(\Theta_\sigma^*(\mathcal{E}))$ .*

**PROOF.** Let us first compute  $\Theta_\sigma^*(\check{c}_\mathbf{x})$ . One has:

$$\begin{aligned} \Theta_\sigma^*(\check{c}_\mathbf{x}) &= \bigcup_{i=1,2,3} \Theta_\sigma^*(\mathbf{x}, i^*) = \bigcup_{\substack{j,p,i,s \\ \sigma(j)=p \cdot i \cdot s}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(s)), j^*) \\ &= \bigcup_{\substack{j,p' \neq \varepsilon, s \neq \varepsilon \\ \sigma(j)=p' \cdot s}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(s)), j^*) \cup \underbrace{\bigcup_{j=1,2,3} (\mathbf{M}_\sigma^{-1} \mathbf{x}, j^*)}_{\check{c}_{\mathbf{M}_\sigma^{-1} \mathbf{x}}}. \end{aligned}$$

and

$$\begin{aligned}
\Theta_\sigma^*(\hat{c}_x) &= \bigcup_{i=1,2,3} \Theta_\sigma^*(\mathbf{x} + \mathbf{e}_i, i^*) = \bigcup_{\substack{j,p,i,s \\ \sigma(j)=p \cdot i \cdot s}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{e}_i + \mathbf{f}(s)), j^*) \\
&= \bigcup_{\substack{j,p,i,s \\ \sigma(j)=p \cdot i \cdot s}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(i \cdot s)), j^*) = \bigcup_{\substack{j,p,s' \neq \varepsilon \\ \sigma(j)=p \cdot s'}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(s')), j^*) \\
&= \bigcup_{\substack{j,p \neq \varepsilon, s' \neq \varepsilon \\ \sigma(j)=p \cdot s'}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(s')), j^*) \cup \bigcup_{j=1,2,3} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(\sigma(j))), j^*) \\
&= \bigcup_{\substack{j,p \neq \varepsilon, s' \neq \varepsilon \\ \sigma(j)=p \cdot s'}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(s')), j^*) \cup \bigcup_{j=1,2,3} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{M}_\sigma \mathbf{e}_j), j^*) \\
&= \bigcup_{\substack{j,p \neq \varepsilon, s' \neq \varepsilon \\ \sigma(j)=p \cdot s'}} (\mathbf{M}_\sigma^{-1}(\mathbf{x} + \mathbf{f}(s')), j^*) \cup \underbrace{\bigcup_{j=1,2,3} (\mathbf{M}_\sigma^{-1} \mathbf{x} + \mathbf{e}_j, j^*)}_{\hat{c}_{\mathbf{M}_\sigma^{-1} \mathbf{x}}},
\end{aligned}$$

since  $\mathbf{e}_i = \mathbf{f}(i)$  and  $\mathbf{f}(\sigma(j)) = \mathbf{M}_\sigma \mathbf{e}_j$ . The desired result easily follows.  $\blacksquare$




Figure 14. If two unions of faces differ by the flip  $\varphi_s$ , then their images by  $\Theta_\sigma^*$  differ by the flip  $\varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}}$  (one has here  $\sigma : 1 \mapsto 13, 2 \mapsto 1, 3 \mapsto 2$ ).

**Lemma 17** *Let  $\mathfrak{S}$  be a stepped surface and  $(\varphi_{\mathbf{x}_n})_{n \in \mathbb{N}}$  be a locally finite sequence of flips such that the sequence  $(\varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{S}))_{n \in \mathbb{N}}$  is convergent in  $\mathcal{G}$ . Then, the sequence of flips  $(\varphi_{\mathbf{M}_\sigma^{-1} \mathbf{v}_n})_{n \in \mathbb{N}}$  is locally finite.*

PROOF. We set  $\mathfrak{S}' = \lim_{n \rightarrow \infty} \varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{S})$ . According to Proposition 10,  $\mathfrak{S}'$  is a stepped surface. Suppose that  $(\varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}_n})_{n \in \mathbb{N}}$  is not locally finite. Let us prove that this implies  $\mathfrak{S}$  is not a stepped surface, which yields a contradiction. We first assume w.l.o.g. that for all  $n \in \mathbb{N}$ , either  $\check{c}_{\mathbf{x}_n}$  or  $\hat{c}_x$  is a subset of  $\varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{S})$ . Since  $(\varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}_n})_{n \in \mathbb{N}}$  is not locally finite, there exists a subsequence  $(\mathbf{y}_n)_{n \in \mathbb{N}}$  of  $(\mathbf{x}_n)_{n \in \mathbb{N}}$ , with  $\sup_n \|\mathbf{y}_n\| = \infty$ , such that:

$$\forall (m, n) \in \mathbb{N}^2, \quad \pi(\mathbf{M}_\sigma^{-1} \mathbf{y}_m) = \pi(\mathbf{M}_\sigma^{-1} \mathbf{y}_n).$$

If we denote by  $\mathbf{u}$  the vector  $\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$ , this is equivalent to say that there exists a sequence  $(\lambda_n) \in \mathbb{Z}^{\mathbb{N}}$ , with  $\sup_n |\lambda_n| = \infty$ , such that:

$$\forall n \in \mathbb{N}, \quad \mathbf{M}_\sigma^{-1}(\mathbf{y}_n - \mathbf{y}_0) = \lambda_n \mathbf{u}.$$

The matrix  $\mathbf{M}_\sigma$  admits nonnegative entries, and at least one positive entry in each row, since  $\det(\mathbf{M}_\sigma) \neq 0$ . Hence the vector  $\mathbf{M}_\sigma \mathbf{u}$  has positive entries.

Moreover, one can assume  $\sup_n \lambda_n = \infty$  (the case  $\inf_n \lambda_n = -\infty$  can be similarly handled). In addition with  $\mathbf{y}_n = \mathbf{y}_0 + \lambda_n \mathbf{M}_\sigma \mathbf{u}$ , where  $(y_{n,1}, y_{n,2}, y_{n,3})$  stands for the entries of  $\mathbf{y}_n$ , this yields:

$$\lim_{n \rightarrow \infty} y_{n,1} = \lim_{n \rightarrow \infty} y_{n,2} = \lim_{n \rightarrow \infty} y_{n,3} = \infty.$$

We have assumed that for all  $n$ ,  $\varphi_{\mathbf{x}_n}$  belongs to the stepped surface  $\varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{S})$ , which is hence above  $\hat{\mathcal{C}}_{\mathbf{x}_n}$ , according to Proposition 8. Let us consider the vertex  $\mathbf{a}_n$  of this stepped surface whose image by  $\pi$  is  $\mathbf{0}$ . This vertex has three identical entries, say,  $\mathbf{a}_n = (a_n, a_n, a_n)$  and is above  $\hat{\mathcal{C}}_{\mathbf{y}_n}$ . Hence,  $a_n \geq \min(y_{n,1}, y_{n,2}, y_{n,3})$ , and therefore,  $\lim_n a_n = \infty$ . Consider now the vertex  $\mathbf{a}_\infty = (a_\infty, a_\infty, a_\infty)$  of  $\mathfrak{S}'$  whose image by  $\pi$  is  $\mathbf{0}$ . For  $n$  large enough,  $\mathbf{a}_\infty$  belongs to  $\varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{S})$  and  $\mathbf{a}_\infty \neq \mathbf{a}_n$ , which yields a contradiction. ■

We are now in a position to prove Theorem 14:

PROOF. Let us consider a stepped surface  $\mathfrak{S}$ . According to Theorem 12, there exist a locally finite sequence of flips  $(\varphi_{\mathbf{x}_n})_{n \in \mathbb{N}}$  and a stepped plane  $\mathfrak{P}$ , such that  $\mathfrak{S}$  can be obtained by performing on  $\mathfrak{P}$  the sequence of flips  $(\varphi_{\mathbf{x}_n})$ :

$$\mathfrak{S} = \lim_{n \rightarrow \infty} \varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{P}).$$

Then, Lemma 15 yields:

$$\Theta_\sigma^*(\mathfrak{S}) = \Theta_\sigma^* \left( \lim_{n \rightarrow \infty} \varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{P}) \right) = \lim_{n \rightarrow \infty} \Theta_\sigma^*(\varphi_{\mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{x}_1}(\mathfrak{P})),$$

and by Lemma 16 one has:

$$\Theta_\sigma^*(\mathfrak{S}) = \lim_{n \rightarrow \infty} \varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}_n} \circ \dots \circ \varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}_1}(\Theta_\sigma^*(\mathfrak{P})).$$

By Theorem 13,  $\Theta_\sigma^*$  maps properly  $\mathfrak{P}$  onto the stepped plane  $\Theta_\sigma^*(\mathfrak{P})$ . The sequence of flips  $(\varphi_{\mathbf{M}_\sigma^{-1} \mathbf{x}_n})_n$  is locally finite by Lemma 17, hence Theorem 12 yields that  $\Theta_\sigma^*(\mathfrak{S})$  is a stepped surface. ■

## References

- [AS03] J.-P. Allouche and J. Shallit. Automatic sequences: Theory and Applications. Cambridge University Press, Cambridge, 2003.
- [AI01] P. Arnoux and S. Ito. Pisot substitutions and Rauzy fractals. *Bull. Belg. Math. Soc. Simon Stevin*, 8(2):181–207, 2001.
- [ABI02] P. Arnoux, V. Berthé, and S. Ito. Discrete planes,  $\mathbb{Z}^2$ -actions, Jacobi-Perron algorithm and substitutions. *Ann. Inst. Fourier (Grenoble)*, 52(2):305–349, 2002.

- [ABJ05] P. Arnoux, V. Berthé, and D. Jamet. Generalized substitutions and stepped surfaces. *Words 2005, 5th International Conference on words (2005)*. Publications du LACIM, 36:101–111, 2005.
- [ABS04] P. Arnoux, V. Berthé, and A. Siegel. Two-dimensional iterated morphisms and discrete planes. *Theoret. Comput. Sci.*, 319(1-3):145–176, 2004.
- [AIS01] P. Arnoux, S. Ito, and Y. Sano. Higher dimensional extensions of substitutions and their dual maps. *J. Anal. Math.*, 83:183–206, 2001.
- [BCK04] V. Brimkov, D. Coeurjolly, and R. Klette. Digital planarity - a review. Technical Report RR2004-24, Laboratoire LIRIS - Université Lumière Lyon 2, may 2004.
- [BV00] V. Berthé and L. Vuillon. Tilings and rotations on the torus: a two-dimensional generalization of sturmian sequences. *Discrete Math.*, 223(1-3):27–53, 2000.
- [Fer05a] T. Fernique. Bidimensional sturmian sequences and substitutions. *DLT'05. LNCS*, 3572:236–247, 2005.
- [Fer05b] T. Fernique. Multidimensional sturmian words and substitutions. *International Journal of Foundations of Computer Science*. Submitted.
- [Fer05c] T. Fernique. Substitutions on multidimensional words. *Words 2005, 5th international conference on words*. Publications du LACIM, 36:223–234, 2005.
- [IO93] S. Ito and M. Ohtsuki. Modified Jacobi-Perron algorithm and generating markov partitions for special hyperbolic toral automorphisms. *Tokyo J. Math.*, 16:441–472, 1993.
- [IO94] S. Ito and M. Ohtsuki. Parallelogram tilings and Jacobi-Perron algorithm. *Tokyo J. Math.*, 17:33–58, 1994.
- [Jam04] D. Jamet. On the Language of Discrete Planes and Surfaces. In *Proceedings of the Tenth International Workshop on Combinatorial Image Analysis*, pages 227-241. Springer-Verlag, 2004.
- [JP05] D. Jamet and G. Paquin, *Discrete surfaces and infinite smooth words*, FPSAC'05-17th Annual International Conference on Formal Power Series and Algebraic Combinatorics, June 20–25, 2005 Taormina, Italy.
- [KR04] R. Klette and A. Rosenfeld, *Digital straightness-A review*, *Discrete Applied Mathematics*, 139:197–230, 2004.
- [Loth02] N. Lothaire, *Algebraic combinatorics on words*, Cambridge University Press, 2002.

- [Rev91] J.-P. Reveillès, *Géométrie discrète, calcul en nombres entiers et algorithmique*, Thèse de doctorat, Université Louis Pasteur, Strasbourg, 1991.
- [Thu89] W. P. Thurston, *Groups, tilings and finite state automata*, Lectures notes distributed in conjunction with the Colloquium Series, in AMS Colloquium lectures, 1989.