

Excursions sur les murs de briques

Khaydar Nurligareev

LIP6, Sorbonne Université

(avec Sergey Kirgizov et Michael Wallner)

Séminaire du GT en Probabilités,
Théorie Ergodique et Systèmes Dynamiques

LMRS, Université de Rouen Normandie

1 décembre, 2025

Short random walks

- d is the dimension,
- $\nu = \frac{d}{2} - 1$,
- $m = \#$ steps,
- A_k is a random step, $|A_k| = 1$.

Short random walks

- d is the dimension,

$$d = 2$$

$$m = 3$$

- $\nu = \frac{d}{2} - 1,$

- $m = \#$ steps,

- A_k is a random step, $|A_k| = 1$.



Short random walks


- d is the dimension,

$$d = 2 \qquad m = 3$$

- $\nu = \frac{d}{2} - 1,$

- $m = \#$ steps,

- A_k is a random step, $|A_k| = 1$.




Short random walks

- d is the dimension,
- $\nu = \frac{d}{2} - 1$,
- $m = \#$ steps,
- A_k is a random step, $|A_k| = 1$.

$$d = 2$$


$$m = 3$$



Short random walks

- d is the dimension,
- $\nu = \frac{d}{2} - 1$,
- $m = \#$ steps,
- A_k is a random step, $|A_k| = 1$.


$d = 2$ $m = 3$



Short random walks

- d is the dimension,
- $\nu = \frac{d}{2} - 1$,
- $m = \#$ steps,
- A_k is a random step, $|A_k| = 1$.

$$d = 2 \quad m = 3$$



Key object: moments $W_m(\nu, n) = \mathbb{E}(|A_1 + \dots + A_m|^n)$

- Fact: for any $m, n \in \mathbb{Z}_{\geq 0}$,
 $W_m(0, 2n)$ and $W_m(1, 2n)$ are integers. Interpretation?

Part I

Planar case, $d = 2$

Matrix form for $d = 2$ ($\nu = 0$). Example: $m = 2$

Fact:

$$W_m(0, 2n) = \sum_{\ell=0}^n [M^{m-1}]_{n\ell}, \quad \text{where } M = \left(\binom{i}{j}^2 \right)_{i,j \geq 0}$$

Example: $m = 2$,

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 4 & 1 & 0 & \cdots \\ 1 & 9 & 9 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Matrix form for $d = 2$ ($\nu = 0$). Example: $m = 2$

Fact:

$$W_m(0, 2n) = \sum_{\ell=0}^n [M^{m-1}]_{n\ell}, \quad \text{where } M = \left(\binom{i}{j}^2 \right)_{i,j \geq 0}$$

Example: $m = 2, \quad W_2(0, 2n) = \binom{2n}{n}$

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 4 & 1 & 0 & \cdots \\ 1 & 9 & 9 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{array}{l} \rightarrow 1 \\ \rightarrow 2 \\ \rightarrow 6 \\ \rightarrow 20 \\ \vdots \end{array}$$

Matrix form for $d = 2$ ($\nu = 0$). Example: $m = 2$

Fact:

$$W_m(0, 2n) = \sum_{\ell=0}^n [M^{m-1}]_{n\ell}, \quad \text{where} \quad M = \left(\binom{i}{j}^2 \right)_{i,j \geq 0}$$

Example: $m = 2, \quad W_2(0, 2n) = \binom{2n}{n}$

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & \dots \\ \color{blue}{1} & \color{blue}{4} & \color{blue}{1} & \color{blue}{0} & \dots \\ 1 & 9 & 9 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{array}{l} \rightarrow 1 \\ \rightarrow 2 \\ \rightarrow \color{blue}{6} \\ \rightarrow 20 \\ \vdots \end{array}$$

$$W_2(0, 4) = 6 :$$

UUDD
UDUD
UDDU
DUUD
DUDU
DDUU

Matrix form for $d = 2$ ($\nu = 0$). Example: $m = 3$

Example: $m = 3$

$$M^2 = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 2 & 1 & 0 & 0 & \cdots \\ 6 & 8 & 1 & 0 & \cdots \\ 20 & 46 & 10 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Matrix form for $d = 2$ ($\nu = 0$). Example: $m = 3$

Example: $m = 3$


$$M^2 = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 2 & 1 & 0 & 0 & \cdots \\ 6 & 8 & 1 & 0 & \cdots \\ 20 & 46 & 10 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{array}{l} \rightarrow 1 \\ \rightarrow 3 \\ \rightarrow 15 \\ \rightarrow 77 \\ \vdots \end{array}$$

Matrix form for $d = 2$ ($\nu = 0$). Example: $m = 3$

Example: $m = 3$

$W_3(0, 4) = 15$:

$$M^2 = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 2 & 1 & 0 & 0 & \cdots \\ 6 & 8 & 1 & 0 & \cdots \\ 20 & 46 & 10 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{matrix} \rightarrow 1 \\ \rightarrow 3 \\ \rightarrow 15 \\ \rightarrow 77 \\ \vdots \end{matrix}$$



RLRL
 RLUD
 RLDU
 RUDL
 RDUL
 ULRD
 UDRL
 DLRU
 DURL
 UUDD
 UDUD
 UDDU
 DUUD
 DUDU
 DDUU

Interpretation for $d = 2$ ($\nu = 0$)

Let $A_k \in \mathbb{C}$, $|A_k| = 1$ ($k = 1, \dots, m$).

$$W_m(0, 2n) = \mathbb{E} |A_1 + \dots + A_m|^{2n}$$

Interpretation for $d = 2$ ($\nu = 0$)

Let $A_k \in \mathbb{C}$, $|A_k| = 1$ ($k = 1, \dots, m$).

$$W_m(0, 2n) = \mathbb{E} |A_1 + \dots + A_m|^{2n}$$

$$\stackrel{(1)}{=} \mathbb{E} \left((A_1 + \dots + A_m) (A_1^{-1} + \dots + A_m^{-1}) \right)^n$$

$$\boxed{1} \quad 1 = |A_k|^2 = A_k \bar{A}_k \quad \Rightarrow \quad A_k^{-1} = \bar{A}_k$$

Interpretation for $d = 2$ ($\nu = 0$)

Let $A_k \in \mathbb{C}$, $|A_k| = 1$ ($k = 1, \dots, m$).

$$W_m(0, 2n) = \mathbb{E} |A_1 + \dots + A_m|^{2n}$$

$$\stackrel{(1)}{=} \mathbb{E} \left((A_1 + \dots + A_m) (A_1^{-1} + \dots + A_m^{-1}) \right)^n$$


$$\stackrel{(2)}{=} [A_1^0 \dots A_m^0] \left((A_1 + \dots + A_m) (A_1^{-1} + \dots + A_m^{-1}) \right)^n$$

$$\boxed{1} \quad 1 = |A_k|^2 = A_k \bar{A}_k \quad \Rightarrow \quad A_k^{-1} = \bar{A}_k$$

$$\boxed{2} \quad \mathbb{E} (A_1 A_2^{-1} A_3 A_1^{-1} A_3 A_2^{-1}) = \mathbb{E} (A_2^{-2} A_3^2) = \mathbb{E} (A_2^{-2}) \cdot \mathbb{E} (A_3^2) = 0$$

Interpretation for $d = 2$ ($\nu = 0$) and $m = 3$



$$\mathbb{E} \left(\left(\begin{array}{ccc} A_1 & + & A_2 & + & A_3 \end{array} \right) \left(\begin{array}{ccc} A_1^{-1} & + & A_2^{-1} & + & A_3^{-1} \end{array} \right) \right)^n$$
$$\begin{array}{cccccc} \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ e_1 & & e_2 & & e_3 & & -e_1 & & -e_2 & & -e_3 \end{array}$$



Interpretation for $d = 2$ ($\nu = 0$) and $m = 3$

$$\mathbb{E} \left(\left(\begin{array}{ccc} A_1 & + & A_2 & + & A_3 \end{array} \right) \left(\begin{array}{ccc} A_1^{-1} & + & A_2^{-1} & + & A_3^{-1} \end{array} \right) \right)^n$$

$$\begin{array}{cccccc} \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ e_1 & & e_2 & & e_3 & & -e_1 & & -e_2 & & -e_3 \end{array}$$




$$A_1 A_2^{-1} A_3 A_1^{-1} A_3 A_2^{-1}$$


Interpretation for $d = 2$ ($\nu = 0$) and $m = 3$

$$\mathbb{E} \left(\left(\begin{array}{ccc} A_1 & + & A_2 & + & A_3 \end{array} \right) \left(\begin{array}{ccc} A_1^{-1} & + & A_2^{-1} & + & A_3^{-1} \end{array} \right) \right)^n$$

$$\begin{array}{cccccc} \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ e_1 & & e_2 & & e_3 & & -e_1 & & -e_2 & & -e_3 \end{array}$$



terms
↓
paths



Interpretation for $d = 2$ ($\nu = 0$) and $m = 3$

$$\mathbb{E} \left(\left(\begin{array}{ccc} A_1 & + & A_2 & + & A_3 \end{array} \right) \left(\begin{array}{ccc} A_1^{-1} & + & A_2^{-1} & + & A_3^{-1} \end{array} \right) \right)^n$$

$\downarrow$
 e_1


$\downarrow$
 e_2

$\downarrow$
 e_3

$\downarrow$
 $-e_1$

$\downarrow$
 $-e_2$


$\downarrow$
 $-e_3$



constant
term




closed
paths



Matrix form revisited for $d = 2$ ($\nu = 0$) and $m = 3$

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions.




Matrix form revisited for $d = 2$ ($\nu = 0$) and $m = 3$

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions.

Let

- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.



Then
$$W_3(0, 2n) = \sum_{k=0}^n \binom{n}{n-k} \binom{n}{n-k} \binom{2k}{k}$$


Matrix form revisited for $d = 2$ ($\nu = 0$) and $m = 3$

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions.

Let

- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.



$$\begin{aligned} \text{Then} \quad W_3(0, 2n) &= \sum_{k=0}^n \binom{n}{n-k} \binom{n}{n-k} \binom{2k}{k} \\ &= \sum_{k=0}^n \binom{n}{k}^2 \sum_{\ell=0}^k \binom{k}{\ell}^2 \end{aligned}$$


Matrix form revisited for $d = 2$ ($\nu = 0$) and $m = 3$

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions.

Let

- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.



$$\begin{aligned} \text{Then} \quad W_3(0, 2n) &= \sum_{k=0}^n \binom{n}{n-k} \binom{n}{n-k} \binom{2k}{k} \\ &= \sum_{k=0}^n \binom{n}{k}^2 \sum_{\ell=0}^k \binom{k}{\ell}^2 \\ &= \sum_{k, \ell=0}^n M_{nk} M_{k\ell} = \sum_{\ell=0}^n [M^2]_{n\ell} \end{aligned}$$

General construction for $d = 2$ ($\nu = 0$)

Paths $\leftrightarrow$ words on $\{U, D, R_1, L_1, \dots, R_{m-2}, L_{m-2}\}$:

- R_s on odd positions,
- L_s on even positions.

Let

- $2n = \# \text{ steps}$,
- $k_0 = \#U = \#D$,
- $k_s = \#R_s = \#L_s$.

$$\begin{aligned} W_m(0, 2n) &= \sum_{k_0 + \dots + k_{m-2} = n} \binom{n}{k_{m-2}}^2 \binom{n-k_{m-2}}{k_{m-3}}^2 \dots \binom{k_1+k_0}{k_0}^2 \binom{2k_0}{k_0} \\ &= \sum_{k_0 + \dots + k_{m-2} = n} \binom{k_0 + \dots + k_{m-2}}{k_0 + \dots + k_{m-3}}^2 \dots \binom{k_0 + k_1}{k_0}^2 \sum_{\ell=0}^n \binom{k_0}{\ell}^2 \\ &= \sum_{\ell, r_1, \dots, r_{m-2}=0}^n M_{nr_{m-2}} \dots M_{r_2 r_1} M_{r_1 \ell} = \sum_{\ell=0}^n [M^{m-1}]_{n\ell} \end{aligned}$$

Summary for $d = 2$ ($\nu = 0$)

- We consider **moments** $W_m(0, 2n) = \mathbb{E}\left(|A_1 + \dots + A_m|^n\right)$,
where $A_k \in \mathbb{C}$, $|A_k| = 1$ ($k = 1, \dots, m$).
- $W_m(0, 2n)$ is the constant term in
$$\left((A_1 + \dots + A_m)(A_1^{-1} + \dots + A_m^{-1})\right)^n.$$
- Thus, $W_m(0, 2n)$ **can be interpreted as** the number of
closed paths of length $2n$ on a specific **m -dimensional lattice**.
- In particular,

$$W_m(0, 2n) = \sum_{\ell=0}^n [M^{m-1}]_{n\ell}, \quad \text{where } M = \left(\binom{i}{j}^2 \right)_{i,j \geq 0}$$


Interlude

Bijection

Brick walls, plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- $2n = \#$ steps,




$\#\{\text{closed paths of size } n\} =$

Brick walls, plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n$$

Brick walls, plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n \binom{n}{n-k}$$



Brick walls, plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n \binom{n}{n-k} \binom{n}{n-k}$$



Brick walls, plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- $2n = \# \text{ steps}$,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n \binom{n}{n-k} \binom{n}{n-k} \binom{2k}{k}$$

R L D L U L R U R D

Brick walls, plane and horizontal half-plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- in each prefix, $\#U \geq \#D$,
- $2n = \#$ steps,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n \binom{n}{n-k} \binom{n}{n-k} C_k$$

R L U L D L R U R D

Brick walls, vertical half-plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $2n = \#$ steps,




$\#\{\text{closed paths of size } n\} =$

Brick walls, vertical half-plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $2n = \#$ steps,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n$$

Brick walls, vertical half-plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $2n = \#$ steps,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.




$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n t_{n+1,k+1}$$

R R L L R L

Brick walls, vertical half-plane

Paths $\leftrightarrow$ words on $\{U, D, R, L\}$:

- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $2n = \#$ steps,
- $k = \#U = \#D$,
- $n - k = \#R = \#L$.



$$\#\{\text{closed paths of size } n\} = \sum_{k=0}^n t_{n+1,k+1} \binom{2k}{k}$$

R D R L U L R U D L

Bijection lemma

The Narayana number

$$t_{n+1, k+1} = \frac{1}{n+1} \binom{n+1}{k} \binom{n+1}{k+1}$$

counts:

words on $\{R, L, O\}$ such that

- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $\#R = \#L = n - k$,
- $\#O = 2k$.

R L R O R O O L O L

Bijection lemma

The Narayana number

$$t_{n+1, k+1} = \frac{1}{n+1} \binom{n+1}{k} \binom{n+1}{k+1}$$

counts:


words on $\{R, L, O\}$ such that

- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $\#R = \#L = n - k$,
- $\#O = 2k$.

R L R O R O O L O L

Dyck paths:

- of size $(n + 1)$,
- with $(k + 1)$ peaks.



Bijection lemma

The Narayana number

$$t_{n+1, k+1} = \frac{1}{n+1} \binom{n+1}{k} \binom{n+1}{k+1} = t_{n+1, n-k+1}$$

counts:

words on $\{R, L, O\}$ such that


- R on odd positions,
- L on even positions,
- in each prefix, $\#R \geq \#L$,
- $\#R = \#L = k$,
- $\#O = 2(n - k)$.

O O R O R O O L O L

$\rightsquigarrow$

Dyck paths:

- of size $(n + 1)$,
- with $(k + 1)$ peaks.



Bijection, part I

- 1 Take a word.

R O O O R O O L O L O O R O R O O L O L

Bijection, part I

- 1 Take a word.
- 2 Identify
 - external R's and L's,

R O O O R O O L O L O O R O R O O L O L

Bijection, part I

- 1 Take a word.
- 2 Identify
 - external R's and L's,
 - external O's.

R O O O R O O L O L O O R O R O O L O L

Bijection, part I

- 1 Take a word.
- 2 Identify
 - external R's and L's,
 - external O's.
- 3 Add one more R and L:
 - R: just before the first external O,
 - L: in the end.

R O O O R O O L O L R O O R O R O O L O L L

Bijection, part I

- 1 Take a word.
- 2 Identify
 - external R's and L's,
 - external O's.
- 3 Add one more R and L:
 - R: just before the first external O,
 - L: in the end.
- 4 Determine subwords $R\alpha_k L$.

R O O O R O O L O L R O O R O R O O L O L L L

$R\alpha_1 L$ $R\alpha_2 L$

Bijection, part I

- 1 Take a word.
- 2 Identify
 - external R's and L's,
 - external O's.
- 3 Add one more R and L:
 - R: just before the first external O,
 - L: in the end.
- 4 Determine subwords $R\alpha_k L$.
- 5 Treat each subword separately.

R O O O R O O L O L R O O R O R O O L O L L L

$R\alpha_1 L$ $R\alpha_2 L$

Bijection, part II

$$\begin{array}{ccccccc}
 R & \underline{O \dots O} & \underline{R\beta_1 L} & \underline{O O \dots O O} & \dots & \underline{O O \dots O O} & \underline{R\beta_m L} & \underline{O \dots O} & L \\
 x & & & 2i_1 & & 2i_{m-1} & & y &
 \end{array}$$

Bijection, part II

$$\begin{array}{ccccccc}
 R & \underline{O \dots O} & \underline{R\beta_1 L} & \underline{O O \dots O O} & \dots & \underline{O O \dots O O} & \underline{R\beta_m L} & \underline{O \dots O} & L \\
 x & & & 2i_1 & & 2i_{m-1} & & y
 \end{array}$$

$$2k = \begin{cases} x - y & \text{if } x > y \\ y - x + 2 & \text{if } x \leq y \end{cases} \quad \downarrow \quad j = \begin{cases} y & \text{if } x > y \\ x - 1 & \text{if } x \leq y \end{cases}$$

$$\begin{array}{ccccccc}
 R & \underline{O O O \dots O O O} & \underline{R\beta_1 L} & \underline{O \dots O} & \dots & \underline{O \dots O} & \underline{R\beta_m L} & \underline{O \dots O} & L \\
 j + 2k + i_1 + \dots + i_{m-1} & & i_1 & & i_{m-1} & & j
 \end{array}$$

Bijection, part II

$$\begin{array}{ccccccc} R & \underline{O \dots O} & \underline{R\beta_1 L} & \underline{O O \dots O O} & \dots & \underline{O O \dots O O} & \underline{R\beta_m L} & \underline{O \dots O} & L \\ x & & & 2i_1 & & 2i_{m-1} & & y \end{array}$$

$$2k = \begin{cases} x - y & \text{if } x > y \\ y - x + 2 & \text{if } x \leq y \end{cases} \quad \downarrow \quad j = \begin{cases} y & \text{if } x > y \\ x - 1 & \text{if } x \leq y \end{cases}$$


$$\begin{array}{ccccccc} R & \underline{O O O \dots O O O} & \underline{R\beta_1 L} & \underline{O \dots O} & \dots & \underline{O \dots O} & \underline{R\beta_m L} & \underline{O \dots O} & L \\ j + 2k + i_1 + \dots + i_{m-1} & & & i_1 & & i_{m-1} & & j \end{array}$$



$$\begin{array}{ccccccc} R & \underline{R R R \dots R R} & \underline{L} & \underline{R\beta_1 L} & \underline{L \dots L} & \dots & \underline{L \dots L} & \underline{R\beta_m L} & \underline{L \dots L} & L \\ j + 2k + i_1 + \dots + i_{m-1} & & & i_1 & & i_{m-1} & & j \end{array}$$


Bijection, example

R O R L O O R L O L R O O R O R O O L O L L



Bijection, example


R O R L O O R L O L R O O R O R O O L O L L



Bijection, example

R O R L O O R L O L R O O R O R O O L O L L


R O O O R L O R L L R O O R O R O O L O L L



Bijection, example


R O R L O O R L O L R O O R O R O O L O L L

R R R L R L L R L L R O O R O R O O L O L L



Bijection, example

R O R L O O R L O L R O O R O R O O L O L L
R R R L R L L R L L L R O O R O R O O L O L L




Bijection, example

R O R L O O R L O L R O O R O R O O L O L L


R R R L R L L R L L R O O R O R O O L O L L

R R R L R L L R L L R L R O R O O L O L L




Bijection, example

R O R L O O R L O L R O O R O R O O L O L L
R R R L R L L R L L R O O R O R O O L O L L
R R R L R L L R L L R R L R O R O O L O L L




Bijection, example

R O R L O O R L O L R O O R O R O O L O L L
R R R L R L L R L L R O O R O R O O L O L L
R R R L R L L R L L R R L R O R O O L O L L
R R R L R L L R L L R R L R O O R O O L L L




Bijection, example

R O R L O O R L O L R O O R O R O O L O L L
R R R L R L L R L L R O O R O R O O L O L L
R R R L R L L R L L R R L R O R O O L O L L
R R R L R L L R L L R R L R R L R O O L L L




Bijection, example

R O R L O O R L O L R O O R O R O O L O L L
R R R L R L L R L L R O O R O R O O L O L L
R R R L R L L R L L R R L R O R O O L O L L
R R R L R L L R L L R R L R R L R O O L L L



Bijection, example

R O R L O O R L O L R O O R O R O O L O L L
R R R L R L L R L L R O O R O R O O L O L L
R R R L R L L R L L R R L R O R O O L O L L
R R R L R L L R L L R R L R R L R O O L L L
R R R L R L L R L L R R L R R L R R L L L



Bijection, example


R O R L O O R L O L R O O R O R O O L O L L

R R R L R L L R L L R O O R O R O O L O L L

R R R L R L L R L L R R L R O R O O L O L L


R R R L R L L R L L R R L R R L R O O L L L

R R R L R L L R L L R R L R R L R R L L L L



Brick walls, quater-planes


Quarter-plane:



$$\# \left\{ \begin{array}{l} \text{closed paths} \\ \text{of size } n \end{array} \right\}$$
$$= \sum_{k=0}^n t_{n+1, k+1} C_k$$


Brick walls, quater-planes

Quarter-plane:



$$\# \left\{ \begin{array}{l} \text{closed paths} \\ \text{of size } n \end{array} \right\} \\ = \sum_{k=0}^n t_{n+1, k+1} C_k$$

Shifted quarter-plane:



$$\# \left\{ \begin{array}{l} \text{closed paths} \\ \text{of size } n + 1 \end{array} \right\} \\ = \sum_{k=0}^n t_{n+1, k+1} C_{k+1}$$

Part II

Four-dimensional case, $d = 4$


Matrix form for $d = 4$ ($\nu = 1$). Example: $m = 2$

Fact:

$$W_m(1, 2n) = \sum_{\ell=0}^n [N^{m-1}]_{n\ell}, \quad \text{where } N = \left(t_{i+1, j+1} \right)_{i, j \geq 0}$$

Here, $t_{i,j}$ are the **Narayana numbers**, i.e.

$$t_{i,j} = \#\{\text{Dyck paths of length } i \text{ with } j \text{ peaks}\}$$



Matrix form for $d = 4$ ($\nu = 1$). Example: $m = 2$

Fact:

$$W_m(1, 2n) = \sum_{\ell=0}^n [N^{m-1}]_{n\ell}, \quad \text{where } N = (t_{i+1,j+1})_{i,j \geq 0}$$

Example: $m = 2$,

$$N = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 3 & 1 & 0 & \cdots \\ 1 & 6 & 6 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Matrix form for $d = 4$ ($\nu = 1$). Example: $m = 2$

Fact:

$$W_m(1, 2n) = \sum_{\ell=0}^n [N^{m-1}]_{n\ell}, \quad \text{where } N = (t_{i+1,j+1})_{i,j \geq 0}$$

Example: $m = 2, \quad W_2(1, 2n) = \frac{1}{n+2} \binom{2n+2}{n+1}$

$$N = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 3 & 1 & 0 & \cdots \\ 1 & 6 & 6 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{array}{l} \rightarrow 1 \\ \rightarrow 2 \\ \rightarrow 5 \\ \rightarrow 14 \\ \vdots \end{array}$$

Matrix form for $d = 4$ ($\nu = 1$). Example: $m = 2$

Fact:

$$W_m(1, 2n) = \sum_{\ell=0}^n [N^{m-1}]_{n\ell}, \quad \text{where } N = (t_{i+1,j+1})_{i,j \geq 0}$$

Example: $m = 2, \quad W_2(1, 2n) = \frac{1}{n+2} \binom{2n+2}{n+1}$

$$N = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ \color{blue}{1} & \color{blue}{3} & \color{blue}{1} & \color{blue}{0} & \cdots \\ 1 & 6 & 6 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad \begin{matrix} \rightarrow 1 \\ \rightarrow 2 \\ \rightarrow \color{blue}{5} \\ \rightarrow 14 \\ \vdots \end{matrix}$$

$W_2(1, 4) = 5 :$



UUUDDD

UUDUDD

UUDDUD


UDUUDD

UDUDUD

Matrix form for $d = 4$ ($\nu = 1$) and $m = 3$

Shifted quarter-plane:

- $2(n+1) = \#$ steps,
- R on **even** positions,
- L on **odd** positions,
- in each prefix, $\#U \geq \#D$,
- in each prefix, $\#R \geq \#L$.




$$\text{Then } \#\{\text{closed paths}\} = \sum_{k=0}^n t_{n+1, k+1} C_{k+1}$$

(here, $k = \#U = \#D$ and $n - k = \#R = \#L$)

Matrix form for $d = 4$ ($\nu = 1$) and $m = 3$

Shifted quarter-plane:

- $2(n+1) = \# \text{ steps}$,
- R on **even** positions,
- L on **odd** positions,
- in each prefix, $\#U \geq \#D$,
- in each prefix, $\#R \geq \#L$.




$$\begin{aligned} \text{Then } \#\{\text{closed paths}\} &= \sum_{k=0}^n t_{n+1, k+1} C_{k+1} \\ &= \sum_{k=0}^n t_{n+1, k+1} \sum_{\ell=0}^k t_{k+1, \ell+1} \end{aligned}$$

(here, $k = \#U = \#D$ and $n - k = \#R = \#L$)

Matrix form for $d = 4$ ($\nu = 1$) and $m = 3$

Shifted quarter-plane:

- $2(n+1) = \# \text{ steps}$,
- R on **even** positions,
- L on **odd** positions,
- in each prefix, $\#U \geq \#D$,
- in each prefix, $\#R \geq \#L$.



$$\begin{aligned}
 \text{Then } \#\{\text{closed paths}\} &= \sum_{k=0}^n t_{n+1, k+1} C_{k+1} \\
 &= \sum_{k=0}^n t_{n+1, k+1} \sum_{\ell=0}^k t_{k+1, \ell+1} \\
 &= \sum_{\ell=0}^n [N^2]_{n\ell} = W_3(1, 2n)
 \end{aligned}$$

(here, $k = \#U = \#D$ and $n - k = \#R = \#L$)

General construction for $d = 4$ ($\nu = 1$)

Paths $\leftrightarrow$ words on $\{U, D, R_1, L_1, \dots, R_{m-2}, L_{m-2}\}$:

- R_s on even positions (after removing all R_t and L_t , $t > s$),
- L_s on odd positions (after removing all R_t and L_t , $t > s$),
- in each prefix, $\#U \geq \#D$ and $\#R_s \geq \#L_s$.

Let

- $2(n+1) = \#$ steps,
- $k_0 = \#U = \#D$,
- $k_s = \#R_s = \#L_s$.

$$\begin{aligned} W_4(1, 2n) &= \sum_{k_0+k_1+k_2=n} t_{n+1, k_1+k_0+1} \cdot t_{n-k_2+1, k_0+1} \cdot C_{k_0+1} \\ &= \sum_{k_0+k_1+k_2=n} t_{n+1, k_1+k_0+1} \cdot t_{k_1+k_0+1, k_0+1} \sum_{\ell=0}^n t_{k_0+1, \ell+1} \\ &= \sum_{\ell, r_1, r_2=0}^n N_{nr_2} N_{r_2 r_1} N_{r_1 \ell} = \sum_{\ell=0}^n [N^3]_{n\ell} \end{aligned}$$

Summary for $d = 4$ ($\nu = 1$)

- We consider **moments** $W_m(1, 2n) = \mathbb{E}(|A_1 + \dots + A_m|^n)$,
where $A_k \in \mathbb{R}^4$, $|A_k| = 1$ ($k = 1, \dots, m$).

- It is known that,

$$W_m(1, 2n) = \sum_{\ell=0}^n [N^{m-1}]_{n\ell}, \quad \text{where } N = (t_{i+1,j+1})_{i,j \geq 0}$$

- Thus, $W_m(1, 2n)$ **can be interpreted as** the number of **closed paths** of length $2(n+1)$ on a specific **m -dimensional lattice**.
- Question. Can we obtain the above result directly?
(one could expect the use of **quaternions**)

Summary for $d = 4$ ($\nu = 1$)

- We consider **moments** $W_m(1, 2n) = \mathbb{E}(|A_1 + \dots + A_m|^n)$,
where $A_k \in \mathbb{R}^4$, $|A_k| = 1$ ($k = 1, \dots, m$).

- It is known that,

$$W_m(1, 2n) = \sum_{\ell=0}^n [N^{m-1}]_{n\ell}, \quad \text{where } N = (t_{i+1,j+1})_{i,j \geq 0}$$

- Thus, $W_m(1, 2n)$ **can be interpreted as** the number of **closed paths** of length $2(n+1)$ on a specific **m -dimensional lattice**.
- Question. Can we obtain the above result directly?
(one could expect the use of **quaternions**)

Thank you for your attention!



Borwein J.M., Straub A., Vignat C.

Densities of short uniform random walks in higher dimensions
J. Math. Anal. Appl., 437(1): pp. 668–707, 2016.